

Analytic and geometric properties of dislocation singularities

Riccardo Scala and Nicolas Van Goethem

Universidade de Lisboa, Faculdade de Ciências,
Departamento de Matemática, CMAFcIO, Alameda
da Universidade, 1749-016 Lisboa, Portugal (rscala@fc.ul.pt;
vangoeth@fc.ul.pt)

(MS Received 22 November 2016; accepted 30 April 2017)

This paper deals with the analysis of the singularities arising from the solutions of the problem $-\text{Curl } F = \mu$, where F is a 3×3 matrix-valued L^p -function ($1 \leq p < 2$) and μ a 3×3 matrix-valued Radon measure concentrated in a closed loop in $\Omega \subset \mathbb{R}^3$, or in a network of such loops (as, for instance, dislocation clusters as observed in single crystals). In particular, we study the topological nature of such dislocation singularities. It is shown that $F = \nabla u$, the absolutely continuous part of the distributional gradient Du of a vector-valued function u of special bounded variation. Furthermore, u can also be seen as a multi-valued field, that is, can be redefined with values in the three-dimensional flat torus \mathbb{T}^3 and hence is Sobolev-regular away from the singular loops. We then analyse the graphs of such maps represented as currents in $\Omega \times \mathbb{T}^3$ and show that their boundaries can be written in term of the measure μ . Readapting some well-known results for Cartesian currents, we recover closure and compactness properties of the class of maps with bounded curl concentrated on dislocation networks. In the spirit of previous work, we finally give some examples of variational problems where such results provide existence of solutions.

Keywords: Cartesian maps; integer-multiplicity currents; torus-valued maps; dislocation singularity; variational problem.

2010 *Mathematics subject classification:* Primary 58A25; 46E40; 32C30; 49Q15; 53C65; 49J45; 74G65

1. Introduction

In this work, we analyse the class of 3×3 matrix-valued fields F defined on an open set $\Omega \subset \mathbb{R}^3$ and having singularities on a 1-dimensional set L . The singularity we are interested in arises from the condition that F is assumed to have a nonzero Curl, which is a finite Radon measure μ concentrated in L , namely

$$-\text{Curl } F = \mu. \tag{1.1}$$

In the simplest case, when the singular set is a closed Lipschitz curve L , it is well known that μ must be a so-called *dislocation density*, that is a measure of

the form

$$\mu = b \otimes \tau \mathcal{H}_{\perp L}^1, \quad (1.2)$$

where b is a fixed vector (constant along L and called the Burgers vector) and τ is the unit tangent vector to L (see lemma 6.1). The term ‘dislocation’ comes from the physics of crystals: it means a line defect where the strain field F is singular. Hence the strain cannot, in general, be associated with a regular deformation since it is constrained to satisfy a condition like (1.1). Equation

$$- \operatorname{Curl} F = b \otimes \tau \mathcal{H}_{\perp L}^1. \quad (1.3)$$

derives from the cut-and-weld model of Volterra, in which the distributional gradient shows a jump part of amplitude b concentrated in the slip plane. Equation (1.3) then follows by taking the curl of its absolutely continuous part (see the model description in [34] and in [33]).

In dislocation theory, the singular set L is always a countable (or finite) union of Lipschitz closed curves, and hence the measure μ is a countable sum of dislocation densities of the form (1.2). Each Burgers vector of the singletons (1.2) is constrained to belong to a countable lattice, that we assume, without loss of generality, to be the group \mathbb{Z}^3 . This setting can be summarized by taking the measure μ in (1.1) in the class of closed integral currents with coefficients in \mathbb{Z}^3 .

By Helmholtz decomposition, any tensor-valued $F \in L^p(\Omega)$ can be written as $F = Du + \operatorname{Curl} V$, with $u \in W^{1,p}(\Omega)$ and V a divergence-free tensor, (see [28, 36]). It is a standard interpretation to call u the displacement field, and V the incompatible field. However, it is immediately observed by (1.1) that the dislocation measure μ depends on V only, hence leaving the displacement outside the picture, at least as far as classically intended as a Sobolev vector-valued map. For this reason, a concept of displacement which keeps into account both u and V at the same time is needed.

The class of vector fields $F : \Omega \rightarrow \mathbb{R}^3$ satisfying condition (1.1) with μ a closed integral current with coefficients in \mathbb{Z}^3 has several measure-theoretic properties that are proved in the present paper. A central aspect is that for every F there exists a 3-dimensional torus-valued map $u : \Omega \rightarrow \mathbb{T}^3$ such that $F = \nabla u$. In some sense, even if the non-vanishing property of $\operatorname{Curl} F$ precludes the existence of a vector-valued function whose gradient is F , we recover such property in the class of torus-valued vector fields. As a consequence, in the spirit of [20], the graph of the map u can be seen as a current and a theory like for Cartesian currents can be developed accordingly. Our main result gives the explicit form of the boundary of the graph associated with the torus-valued deformation in terms of the measure μ (the dislocation density in the specific application). Other forms of this result are also given, as well as a discussion on its consequence for a variational approach to dislocations. Let us emphasize that the use of Cartesian currents and the powerful Federer-Fleming closure theorem was first considered in [33] in order to reconstruct the strain field for a given dislocation density, assumed to be known beforehand. In the present paper, we would like to go one step further, and without the restrictive working assumptions on the geometry of the dislocation lines of our previous work

[37], to find the optimal dislocation density compatible in the far field with a given dislocation density prescribed at the boundary.

1.1. Line-like singularities in Continuum mechanics

A dislocation (or disclination) is a line defect in a crystal related to the jump of the displacement or the rotation tensor, itself provoked by the motion of atoms and atom layers at an underlying microscopic scale. At the mesoscopic scale, where the medium is considered as continuous, such a line represents a singularity for model fields such as stress, strain, displacement, and rotation. Furthermore, in single crystals these lines might appear as isolated loops as well as complex networks and there exist mutual interactions such that two lines might join together or (partially) annihilate, or be set apart in many smaller loops. All these phenomena should be understood as mathematical objects which must be described in a geometrically unified way. To achieve this goal, the theory of currents has shown as extremely appropriate, as described in [37]. Employing the tool of integral currents to describe dislocations has been used by several authors in recent years, see, for instance, [25], [24]. It should be emphasized that integral currents and Cartesian currents also allow one to handle singular problems in Continuum mechanics by means of the notions of distributional Jacobian and cofactors of the deformation tensor. Pioneer works and fundamental results about this topic can be found in [32]. In the recent years, this formalism has been developed for elasticity and, in general, for Continuum mechanics, as in, for example, [21, 30, 31] or [2, 3, 23]. About their specific use in problems involving dislocations, see [33, 37]. About their use in problems involving other kinds of loop discontinuities, the Ginzburg and Landau problem has provided numerous key contributions, as for example, in [1, 7, 35]. Indeed, the mathematical issues addressed in the latter works very often appear in a similar way in the study of defects in solids, which therefore share with Ginzburg-Landau models several theoretical concepts, among which field multiple-valuedness and vorticity. Let us also mention functions of higher-order variation [26] (considered in [37] for dislocation problems), or strings [10], as some of the many mathematical objects arisen from the study of singular loops in PDEs. It should be stressed that one of the features of these problems is that the model fields have either values in a manifold (for instance, in the unit sphere), or are multiple-valued. Multivaluedness is a challenging issue since the pioneer works of Almgren [4], both in Physics problems such as liquid crystals, or from a purely mathematical perspective [8, 16], where often, multivaluedness is related to singularities, as in [11].

Sometimes, dislocations are considered in a unified way with fractures, as for example, in [30], where ‘fracture’ is intended as a non-negative co-dimension singularity, that is, classical fractures have co-dimension 1, whereas dislocations have co-dimension 2, and point defects co-dimension 3. However, dislocation singularities show several specific issues, and the intrinsic mathematical difficulties generated by dislocations are fundamentally different from those encountered in the mathematical modelling of fracture mechanics. Indeed,

- (i) dislocations are \mathcal{H}^1 -singularities in \mathbb{R}^3 and hence show no jump set in the classical sense;
- (ii) dislocations are free to mutually annihilate, recombine, split, spontaneously appear, and hence form complex geometrical structures, without any law such as irreversibility;
- (iii) there is no natural reference configuration and hence intrinsic approaches must be preferred. In particular, the displacement is not an appropriate model variable *in the classical sense*, as opposed to most of solid mechanics problems;
- (iv) the stress and strain fields are not square-integrable: thus the less tractable L^p spaces with $1 \leq p < 2$ must be considered;
- (v) bounds on the model fields are given in terms of the curl and the divergence, in place of the full gradient;
- (vi) these curl and divergence are found in measure spaces instead of Sobolev spaces.

In the language of currents, the measure (1.2) is denoted as $b \otimes \mathcal{L}$, with \mathcal{L} standing for an integral 1-current, whose support is the curve L , as described in [37]. In the language of Physics, it is simply the (transpose of) the dislocation density $\Lambda_L := \tau \otimes b\delta_L$ (see [38]). The tensor-valued functions satisfying the differential constraint (1.1) will be denoted by F to mean the deformation tensor of finite elasticity. Because of this constraint, F belongs to $L^p(\Omega)$ with $1 \leq p < 2$. It is a classical procedure to also consider the determinant and cofactors of F as belonging to the same space since J. Ball results on existence results for polyconvex functions [6, 12]. Such an issue will be addressed in a forthcoming work, as a natural consequence of the analysis of the present paper.

1.2. Scope and structure of the work

In our approach, we follow Almgren and Lieb's understanding of singularities through the theory of Cartesian currents by means of a tractable notion of multi-valuedness [5]. As a first step, our aim is to analyse the fields satisfying the constraint (1.1), showing that an unambiguous concept of displacement can be defined. As a second step, we want to study some measure-theoretic properties of such displacements, proving a series of important closure results. More specifically, it will be shown that

- for all F satisfying (1.1) there exists a map $u \in W^{1,p}(\Omega, \mathbb{T}^3)$, with $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ the flat torus, such that $F = \nabla u$. Moreover, there exists a map $\tilde{u} \in SBV^p(\Omega, \mathbb{R}^3)$ such that $F = \nabla^a \tilde{u}$, with $\nabla^a \tilde{u}$ the absolutely continuous part of the distributional gradient $D\tilde{u}$.
- $\partial \mathcal{G}_u$, the boundary of the current supported by the graph of u , is 'equal' (in a sense that will be made precise) to the dislocation density Λ_L . Moreover, $\Lambda_L^T = -\text{Curl } \nabla u$.

The first item states that it is possible to construct a real vector-valued displacement field \tilde{u} , which has a constant jump b on a fixed (though arbitrary) surface S enclosed by \mathcal{L} . Furthermore, if we quotient the target space \mathbb{R}^3 by \mathbb{Z}^3 , the jump on the (arbitrary) surface can be neglected. In other words, the displacement as a Euclidean-valued field is multiple-valued, since its circulation around dislocation lines is nonzero and is always an integer multiple of b . As an alternative, working with torus-valued maps also allows us to remove this ambiguity. Torus-valued maps are therefore alternatives to Euclidean maps appropriate to mathematically model dislocation singularities. Similarly, it will be convenient to introduce periodic k -forms, from which torus-valued currents are defined by duality since these forms appear as more suitable in this setting. Note that such multivalued displacement fields have been used by Physicists for a long time (see, e.g., [27]), but without the necessary mathematical rigor, which we believe is worth introducing to seek a profound understanding of the intrinsic difficulties generated by dislocation singularities. It should be mentioned that considering the strain F associated with a dislocation measure as the approximate gradient of an SBV -map had been considered before as a heuristic picture in [15, 33]. To our knowledge, the present work is the first to rigorously address this identification.

As for the last item, with the language of currents and in dislocation terms, our main result basically states that the dislocation density is the boundary of the graph of a displacement field u which has values in the three-dimensional torus, viz.,

$$\partial\mathcal{G}_u(\omega) = \mathcal{L} \otimes b(\varphi^\omega),$$

where ω is any 2-form, and φ^ω is a test function depending linearly on ω (see theorem 6.4). Starting from this result it is possible to write the explicit expression of the boundary of the graph of $w = v + u$, where v plays the role of a smooth perturbation of the function u . Furthermore, we give the explicit form of the boundary of the graph of the vector-valued function \tilde{u} , which indeed coincides with the torus-valued u , up to an additional measure concentrated in the jump set of \tilde{u} (theorem 5.1). This is our main result second form, which might be useful in models involving an energy term related to the jump surface, whose complete analysis, though, is left for future works.

Finally, application examples are given: two variational problems are proposed and solved in § 6.5. In particular, in the present work, we state some partial results in the direction of generalizing the minimization problems considered in [37]. Specifically, we consider energy densities which also depend on $\operatorname{div} F$ and, relying on Helmholtz decomposition $F = Du + F^0$ with $\operatorname{div} F^0 = 0$, we consider the cases where (i) $\operatorname{div} F = 0$, and (ii) $\operatorname{div} F \in L^q(\Omega)$ for some appropriate exponent q . Obviously, these results must be considered as a milestone toward more general results to appear in future works [40, 41]. Variational problems of this kind have been already treated by several authors but without the generality of unfixed dislocation loops. The work closest to our approach of minimizing an elastic energy in the presence of dislocations was proposed in the pioneer paper [33], where the authors enriched the theory by presenting an alternative to the core-radius approach in linearized elasto-plasticity theory. This alternative formulation is more appropriate in situations where large strains are present and it is not necessary to introduce a core radius, since the integrability assumption $F \in L^p$, with $p < 2$ is compatible

with unbounded gradients at the dislocation line. Also close to our approach, let us mention [13], where the study is restricted to 2d setting, [14], for an analysis of a surface energy alone and where only the compactness of 1-currents is needed (as opposed to our approach of considering simultaneously the 1-current and its associated strain F), and [15] considering the 3d case, though with loops that are still assumed to be known beforehand.

This paper is structured as follows. In § 2, we recall the basis for currents and the theory of graphs of torus-valued maps, and prove some important preliminary results, whose counterparts in the Euclidean-valued case are classical. Then, the crucial pointwise and distributional properties of torus-valued harmonic maps are derived in § 3. The main results are presented in two forms, the first in § 4 holding for torus-valued maps (see theorem 4.1), and the second one valid for Euclidean maps is given in § 5, with a proof in Appendix A. In § 6, we show how the results obtained so far can be applied to the theory of dislocations, whereas in § 6.5, we provide two examples of variational problems which can be solved by virtue of our results.

2. Preliminary results

2.1. Generalities about currents and graphs

For all integers $n > 0$ and $k \geq 0$ with $k \leq n$, we denote by $\Lambda_k \mathbb{R}^n$ the space of k -vectors and by $\Lambda^k \mathbb{R}^n$ the space of k -covectors. Let α be a multi-index, that is, an ordered (increasing) subset of $\{1, 2, \dots, n\}$. We denote by $|\alpha|$ the cardinality (or length) of α , and we denote by $\bar{\alpha}$ the complementary set of α , that is, the multi-index given by the ordered set $\{1, 2, \dots, n\} \setminus \alpha$.

For a $n \times n$ matrix A with real entries and for α and β multi-indices such that $|\alpha| + |\beta| = n$, $M_{\bar{\alpha}}^{\beta}(A)$ will denote the determinant of the submatrix of A obtained by erasing the i -th columns and the j -th rows, for all $i \in \alpha$ and $j \in \bar{\beta}$. Moreover, symbol $M(A)$ will denote the n -vector in $\Lambda_n \mathbb{R}^{2n}$ given by

$$M(A) := \sum_{|\alpha|+|\beta|=n} \sigma(\alpha, \bar{\alpha}) M_{\bar{\alpha}}^{\beta}(A) e_{\alpha} \wedge \varepsilon_{\beta},$$

where $\{e_i, \varepsilon_i\}_{i \leq n}$ is the Euclidean basis of \mathbb{R}^{2n} , and $\sigma(\alpha, \bar{\alpha})$ is the sign of the permutation $(\alpha, \bar{\alpha})$. Accordingly, define

$$|M(A)| := \left(1 + \sum_{\substack{|\alpha|+|\beta|=n \\ |\beta|>0}} |M_{\bar{\alpha}}^{\beta}(A)|^2 \right)^{1/2}.$$

For a matrix $A \in \mathbb{R}^{3 \times 3}$, the symbols $\text{adj } A$ and $\det A$ denote the adjunct, that is, the transpose of the matrix of the cofactors of A , and the determinant of A ,

respectively. Explicitly,

$$M_j^i(A) = A_{ij}, \quad M_j^{\bar{i}}(A) = (\text{cof } A)_{ij} = (\text{adj } A)_{ji} \quad M_{\{1,2,3\}}^{\{1,2,3\}}(A) = \det A,$$

for $i, j = 1, 2, 3$. Moreover,

$$|M(A)| = \left(1 + \sum_{i,j} A_{ij}^2 + \sum_{i,j} \text{cof}(A)_{ij}^2 + \det(A)^2 \right)^{1/2}. \tag{2.1}$$

Let $\Omega \subset \mathbb{R}^n$ be an open set, then $\mathcal{D}^k(\Omega)$ denotes the topological vector space of compactly supported smooth k -forms in Ω , that is, the topological vector space of compactly supported and smooth maps in Ω with values in $\Lambda^k \mathbb{R}^n$. The dual space of $\mathcal{D}^k(\Omega)$, denoted by $\mathcal{D}_k(\Omega)$, is known as the space of k -currents in Ω . A weak convergence in $\mathcal{D}_k(\Omega)$ is defined by saying that $\mathcal{T}_h \rightharpoonup \mathcal{T}$ as currents if for all $\omega \in \mathcal{D}^k(\Omega)$ we have $\mathcal{T}_h(\omega) \rightarrow \mathcal{T}(\omega)$. The boundary of a current $\mathcal{T} \in \mathcal{D}_k(\Omega)$ is the current $\partial\mathcal{T} \in \mathcal{D}_{k-1}(\Omega)$ defined as

$$\partial\mathcal{T}(\omega) = \mathcal{T}(d\omega), \quad \text{for all } \omega \in \mathcal{D}^{k-1}(\Omega).$$

A current which has null boundary (as any current in $\mathcal{D}_0(\Omega)$) is said closed. For all $\mathcal{T} \in \mathcal{D}_k(\Omega)$ the mass of \mathcal{T} is the number $M(\mathcal{T}) \in [0, +\infty]$ defined as

$$M(\mathcal{T}) := \sup_{\omega \in \mathcal{D}^k(\Omega), |\omega| \leq 1} \mathcal{T}(\omega).$$

If $M(\mathcal{T}) < +\infty$ then \mathcal{T} turns out to be a Borel measure in $\mathcal{M}_b(\Omega, \Lambda^k \mathbb{R}^n)$, and its mass coincides with $M(\mathcal{T})$. Moreover, the mass is lower semicontinuous with respect to the weak topology in $\mathcal{D}_k(\Omega)$. Indeed if $\limsup_{h \rightarrow \infty} M(\mathcal{T}_h) < +\infty$ and $\mathcal{T}_h \rightharpoonup \mathcal{T}$ then we also find that \mathcal{T} is a Borel measure and $\mathcal{T}_h \rightharpoonup \mathcal{T}$ weakly in $\mathcal{M}_b(\Omega, \Lambda^k \mathbb{R}^n)$, so that the lower-semicontinuity of the mass follows from the lower-semicontinuity of the mass in $\mathcal{M}_b(\Omega, \Lambda^k \mathbb{R}^n)$. We also define the quantity

$$N(\mathcal{T}) := M(\mathcal{T}) + M(\partial\mathcal{T}),$$

for every $\mathcal{T} \in \mathcal{D}_k(\Omega)$.

Let $U \subset \mathbb{R}^n$ and $V \subset \mathbb{R}^m$ be open sets and $F : U \rightarrow V$ be a smooth map. Then the *push-forward* of a current $\mathcal{T} \in \mathcal{D}_k(U)$ by F is defined as

$$F_{\#}\mathcal{T}(\omega) := \mathcal{T}(\zeta F^{\sharp}\omega) \quad \text{for } \omega \in \mathcal{D}^k(V),$$

where $F^{\sharp}\omega$ is the standard pull-back of ω and ζ is any \mathcal{C}^∞ function that is equal to 1 on $\text{spt}\mathcal{T} \cap \text{spt}F^{\sharp}\omega$. It turns out that $F_{\#}\mathcal{T} \in \mathcal{D}_k(V)$ does not depend on ζ and satisfies

$$\partial F_{\#}\mathcal{T} = F_{\#}\partial\mathcal{T}. \tag{2.2}$$

Let $0 \leq k \leq n$ and let $S \subset \mathbb{R}^n$ be a \mathcal{H}^k -rectifiable set with approximate tangent space $T_x S$. If $\tau : S \rightarrow \Lambda_k(\mathbb{R}^n)$ and $\theta : S \rightarrow \mathbb{R}$ are \mathcal{H}^k -integrable functions with

$\tau(x) \in T_x S$ a simple unit k -vector for \mathcal{H}^k -a.e. $x \in S$, then we can define the current \mathcal{T} as

$$\mathcal{T}(\omega) = \int_S \langle \omega(x), \tau(x) \rangle \theta(x) d\mathcal{H}^k(x) \quad \text{for } \omega \in \mathcal{D}^k(\Omega). \tag{2.3}$$

Every current for which there exists S , τ , and θ as before is said to be *rectifiable*. If also its boundary $\partial\mathcal{T}$ is rectifiable, then we adopt the following notation

$$\mathcal{T} \equiv \{S, \tau, \theta\}. \tag{2.4}$$

The current $\mathcal{T} \in \mathcal{D}_k(\Omega)$ is *rectifiable with integer multiplicity* if it is rectifiable, has rectifiable boundary, and the function θ in (2.3) is integer-valued. A integer-multiplicity current \mathcal{T} such that $N(\mathcal{T}) < \infty$ is said an *integral current*.

The simplest example of integer-multiplicity current is the integration over a set $U \subset \mathbb{R}^n$. This is denoted by $\llbracket U \rrbracket \in \mathcal{D}_n(\mathbb{R}^n)$ and is defined as

$$\llbracket U \rrbracket(\omega) = \int_U \langle \tau, \omega(x) \rangle dx \quad \text{for all } \omega \in \mathcal{D}^n(\mathbb{R}^n),$$

where $\tau := e_1 \wedge \dots \wedge e_n$ the standard orientating vector of \mathbb{R}^n .

An integer-multiplicity current $\mathcal{T} \in \mathcal{D}_M(\mathbb{R}^n)$ is said *indecomposable* if there exists no integral current \mathcal{R} such that $\mathcal{R} \neq 0 \neq \mathcal{T} - \mathcal{R}$ and

$$N(\mathcal{T}) = N(\mathcal{R}) + N(\mathcal{T} - \mathcal{R}).$$

The following theorem provides a decomposition property of every integral current and the structure characterization of integer-multiplicity indecomposable 1-currents (see [17, § 4.2.25]).

THEOREM 2.1. *For every integral current \mathcal{T} there exists a sequence of indecomposable integral currents \mathcal{T}_i such that*

$$\mathcal{T} = \sum_i \mathcal{T}_i \quad \text{and} \quad N(\mathcal{T}) = \sum_i N(\mathcal{T}_i).$$

Suppose \mathcal{T} is an indecomposable integer multiplicity 1-current on \mathbb{R}^n . Then there exists a Lipschitz function $f : \mathbb{R} \rightarrow \mathbb{R}^n$ with $\text{Lip}(f) \leq 1$ such that

$$f \llcorner [0, M(\mathcal{T})) \text{ is injective and } \mathcal{T} = f \# \llbracket [0, M(\mathcal{T})) \rrbracket.$$

Moreover, $\partial\mathcal{T} = 0$ if and only if $f(0) = f(M(\mathcal{T}))$.

Let us consider the space $\Omega \times \mathbb{R}^3$. We will use the Euclidean coordinates $x = (x_1, x_2, x_3)$ for $x \in \Omega$ and $y = (y_1, y_2, y_3)$ for $y \in \mathbb{R}^3$. Every 3-form $\omega \in \mathcal{D}^3(\Omega \times \mathbb{R}^3)$ can be decomposed as $\omega = \sum \omega_{\alpha\beta} dx^\alpha \wedge dy^\beta$, with $\omega_{\alpha\beta} \in C_c^\infty(\Omega \times \mathbb{R}^3)$, where the sum is computed over all multi-indices α and β such that $|\alpha| + |\beta| = 3$.

For $1 \leq p < +\infty$, we define

$$\mathcal{A}_p(\Omega, \mathbb{R}^3) := \{u \in L^p(\Omega, \mathbb{R}^3) : u \text{ is approximately differentiable a.e. and}$$

$$M_\alpha^\beta(Du) \in L^p(\Omega) \text{ for all } |\alpha| + |\beta| = 3\}.$$

A weak convergence is defined on $\mathcal{A}_p(\Omega, \mathbb{R}^3)$. We say that the sequence $u_k \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$ converges to $u \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$ weakly in $\mathcal{A}_p(\Omega, \mathbb{R}^3)$ if $u_k \rightharpoonup u$ weakly in

$L^p(\Omega, \mathbb{R}^3)$ and $M_{\bar{\alpha}}^{\beta}(Du_k) \rightharpoonup M_{\bar{\alpha}}^{\beta}(Du)$ weakly in $L^p(\Omega)$ for all multi-indices α and β with $|\alpha| + |\beta| = 3$ (see [20, § 3.3.3]). If $u \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$ then \mathcal{G}_u , the current carried by the graph of u , is defined as follows:

$$\mathcal{G}_u := (Id \times u)_{\#} \llbracket \Omega \rrbracket, \tag{2.5}$$

where $Id \times u : \mathbb{R}^3 \rightarrow \mathbb{R}^3 \times \mathbb{R}^3$ is given by $(Id \times u)(x) = (x, u(x))$, viz.,

$$\mathcal{G}_u(\omega) = \int_{\Omega} \sigma(\alpha, \bar{\alpha}) \omega_{\alpha\beta}(x, u(x)) M_{\bar{\alpha}}^{\beta}(Du(x)) dx, \tag{2.6}$$

for all $\omega = \omega_{\alpha\beta} dx^{\alpha} \wedge dy^{\beta} \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$. Moreover, if $u \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$ then \mathcal{G}_u turns out to be a 3-integer-multiplicity current in $\Omega \times \mathbb{R}^3$.

LEMMA 2.2. *Let $u_k, u \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$ be such that $u_k \rightharpoonup u$ weakly in $\mathcal{A}_p(\Omega, \mathbb{R}^3)$, then $\mathcal{G}_{u_k} \rightharpoonup \mathcal{G}_u$ as currents.*

Proof. This is a straightforward consequence of formula (2.6). □

The boundary of the current \mathcal{G}_u is, by definition, the 2-current in $\Omega \times \mathbb{R}^3$ given by $\partial \mathcal{G}_u(\omega) := \mathcal{G}_u(d\omega)$ for all 2-forms $\omega \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$. As soon as $u|_U$, the restriction of u to a smooth open subset U of Ω , is smooth in U , then $\mathcal{G}_{u|_U}$ is a current with rectifiable boundary given by

$$\partial \mathcal{G}_{u|_U} = \mathcal{G}_{u|_{\partial U}} := (Id \times u)_{\#} \llbracket \partial U \rrbracket.$$

We can also employ Stokes formula to find an explicit formula for $\partial \mathcal{G}_{u|_{\partial U}}$, that is,

$$\mathcal{G}_{u|_U}(d\omega) = \partial \mathcal{G}_{u|_U}(\omega) = \int_{\partial U} \omega_{\alpha\beta}(x, u(x)) \left\langle dx^{\alpha} \wedge dy^{\beta}, \frac{\partial \Psi}{\partial \tau_1} \wedge \frac{\partial \Psi}{\partial \tau_2} \right\rangle d\mathcal{H}^2(x), \tag{2.7}$$

for all 2-form $\omega = \omega_{\alpha\beta} dx^{\alpha} \wedge dy^{\beta} \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$, and where $\Psi = Id \times u$. This can be seen as follows: by definition of push-forward of a current, $\mathcal{G}_{u|_{\partial U}}(\omega) = \llbracket \partial U \rrbracket((Id \times u)_{\#}(\omega)) = \int_{\partial U} \langle (Id \times u)_{\#} \omega, \tau_1 \wedge \tau_2 \rangle d\mathcal{H}^2 = \int_{\partial U} \langle \omega, ((\partial \Psi)/(\partial \tau_1)) \wedge ((\partial \Psi)/(\partial \tau_2)) \rangle d\mathcal{H}^2$ where $\tau_1 \wedge \tau_2$ is a volume form for ∂U . Now by Stokes Theorem this is equal to (recall that one can interchange the exterior differentiation and pull-back operations) $\int_U \langle d(Id \times u)_{\#} \omega, e_1 \wedge e_2 \wedge e_3 \rangle d\mathcal{H}^3 = \int_U \langle (Id \times u)_{\#} d\omega, e_1 \wedge e_2 \wedge e_3 \rangle d\mathcal{H}^3 = \mathcal{G}_{u|_U}(d\omega) = \partial \mathcal{G}_{u|_U}(\omega)$ (see [29] for details).

The class of Cartesian maps is the subspace of $\mathcal{A}_1(\Omega, \mathbb{R}^3)$ defined as

$$\text{Cart}^1(\Omega, \mathbb{R}^3) := \{u \in \mathcal{A}_1(\Omega, \mathbb{R}^3) : \partial \mathcal{G}_u = 0\}. \tag{2.8}$$

If $u \in W^{1,p}(\Omega, \mathbb{R}^3)$ with $p > 3$, then it is easy to see that $u \in \text{Cart}^1(\Omega, \mathbb{R}^3)$. See [19, § 3.2.2] for details.

THEOREM 2.3. *Let $p > 1$. Let u_k be a sequence in $\mathcal{A}_p(\Omega, \mathbb{R}^3)$ such that $u_k \rightarrow u$ strongly in $L^p(\Omega, \mathbb{R}^3)$ and suppose that there exist functions $v_{\alpha}^{\beta} \in L^p(\Omega)$ such that $M_{\bar{\alpha}}^{\beta}(Du_k) \rightharpoonup v_{\alpha}^{\beta}$ for all multi-indices α and β with $|\alpha| + |\beta| = 3$. If*

$$M(\partial \mathcal{G}_{u_k}) < C < +\infty \tag{2.9}$$

for all $k > 0$, then $u \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$ and $v_{\alpha}^{\beta} = M_{\bar{\alpha}}^{\beta}(Du)$.

This is proved in theorem 2 of [19, § 3.3.2]. Since Cartesian maps obviously satisfy condition (2.9), this Theorem applies to this case. Moreover, if $u_k \in \text{Cart}^1(\Omega, \mathbb{R}^3)$ then we also have $u \in \text{Cart}^1(\Omega, \mathbb{R}^3)$.

2.2. Torus-valued maps and currents

For simplicity of exposition, we describe the 3-dimensional case, but all the definitions and statements of this section can be established for the general n -dimensional case. We introduce the torus $\mathbb{T} \cong \mathbb{R}/\sim$, where \sim denotes the equivalent relation given by $a \sim b$ iff $a - b \in 2\pi\mathbb{Z}$. We also denote $\mathbb{T} \times \mathbb{T} \times \mathbb{T}$ by \mathbb{T}^3 . Now we will consider graphs of maps $u : \Omega \rightarrow \mathbb{T}^3$. These turn out to be 3-rectifiable currents in $\Omega \times \mathbb{T}^3$. Note that the space of 3-forms in $\Omega \times \mathbb{T}^3$, that is, $\mathcal{D}^3(\Omega \times \mathbb{T}^3)$, is exactly the space of 3-forms in $\Omega \times \mathbb{R}^3$ that have coefficients which are smooth and 2π -periodic (with all their derivatives) in the last three variables (actually, they do not have compact support). As a consequence, if \mathcal{T} is a 3-current in $\mathcal{D}_3(\Omega \times \mathbb{R}^3)$ with compact support in $\bar{\Omega} \times \mathbb{R}^3$, then the current

$$T(\mathcal{T}) := \mathcal{T} \llcorner_{\mathcal{D}^3(\Omega \times \mathbb{T}^3)}, \tag{2.10}$$

is well-defined and $T(\mathcal{T}) \in \mathcal{D}_3(\Omega \times \mathbb{T}^3)$. Moreover, since in general smooth functions in $\Omega \times \mathbb{R}^3$ are not periodic in the last three variables, it turns out that $M(T(\mathcal{T})) \leq M(\mathcal{T})$.

Let $u \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$, then we define $T(u) : \Omega \rightarrow \mathbb{T}^3$ by means of the standard projection $\pi_T : \mathbb{R} \rightarrow \mathbb{T}$, that is, $T(u_i) := \pi_T(u_i)$ for $i = 1, 2, 3$. It is easily seen that \mathbb{T} being locally isomorphic to \mathbb{R} , $T(u)$ is almost everywhere approximately differentiable with the same approximate derivatives of u . As a consequence, $\mathcal{G}_{T(u)}$ is a 3-rectifiable current in $\Omega \times \mathbb{T}^3$. It is also easy to see that in such a case $\mathcal{G}_{T(u)} = T(\mathcal{G}_u)$. This fundamental identity follows from the fact that the approximate differentials of u and $T(u)$ coincide almost everywhere, and from (2.6), where we use that if ω is 2π -periodic in the second variable, then $\omega(x, u(x)) = \omega(x, T(u(x)))$. We introduce the space $\mathcal{A}_p(\Omega, \mathbb{T}^3)$ as follows:

DEFINITION 2.4.

$$\begin{aligned} \mathcal{A}_p(\Omega, \mathbb{T}^3) := \{ & u \in L^1(\Omega, \mathbb{T}^3) : u \text{ is approximately differentiable a.e. on } \Omega, \quad \text{and} \\ & M_\alpha^\beta(Du) \in L^p(\Omega) \quad \text{for all } |\alpha| + |\beta| = 3\}, \end{aligned} \tag{2.11}$$

where $L^1(\Omega, \mathbb{T}^3)$ means the space of measurable functions $u : \Omega \rightarrow \mathbb{T}^3$.

With this definition, we see that for all $u \in \mathcal{A}_p(\Omega, \mathbb{T}^3)$ the graph \mathcal{G}_u is well defined as a 3-rectifiable current. A consequence of the fact that the mass of a current does not increase when we compose with T is that, if there exists $\bar{u} \in \mathcal{A}_p(\Omega, \mathbb{R}^3)$ such that $T(\bar{u}) = u$ and $\mathcal{G}_{\bar{u}}$ is an integral current, then \mathcal{G}_u is an integral current. Note that it might happen that such \bar{u} does exist with $\partial\mathcal{G}_{\bar{u}}$ unbounded, whereas $M(\partial\mathcal{G}_u) < \infty$.

Theorem 2.3, being a consequence of the compactness theorem for integral currents, straightforwardly applies also to the case of maps with values in \mathbb{T}^3 . Also, lemma 2.2 readily applies to the case of maps with value in \mathbb{T}^3 .

LEMMA 2.5. Let $u_k, u \in \mathcal{A}_p(\Omega, \mathbb{T}^3)$ be such that $u_k \rightharpoonup u$ weakly in $\mathcal{A}_p(\Omega, \mathbb{T}^3)$, then $\mathcal{G}_{u_k} \rightharpoonup \mathcal{G}_u$ as currents.

Proof. This is again a consequence of formula (2.6) and the fact that currents in $\mathcal{D}_3(\Omega, \mathbb{T}^3)$ belong also to $\mathcal{D}_3(\Omega, \mathbb{R}^3)$. \square

3. Harmonic maps with prescribed jump on a surface

In this section, we study some regularity results of harmonic maps in \mathbb{R}^3 with prescribed jump on a given surface. This is the first step to investigate the geometric properties of the deformation gradients satisfying equation (1.3). Following the model of Volterra (see [33, 34]), if a dislocation L is the boundary of a surface S , then the displacement can be seen as a function with a fixed jump on the surface, with amplitude the constant Burgers vector associated with L . In this work, we will often need to make the following hypothesis:

ASSUMPTION 3.1. We consider a closed Lipschitz curve L , which is assumed simple, and thus non self-intersecting. Moreover, we assume that S is a bounded Lipschitz simple surface whose boundary is L , that is, S is non self-intersecting and is homeomorphic to the disk $B_1 \subset \mathbb{R}^2$.

LEMMA 3.2. Let L be a Lipschitz closed curve in \mathbb{R}^3 and S a bounded Lipschitz surface with boundary L and unit normal N (satisfying assumption 3.1). Let $b \in \mathbb{R}$. A solution of

$$\begin{cases} \Delta u = 0 & \text{in } \mathbb{R}^3 \setminus S \\ [u] := u^+ - u^- = b & \text{on } S \\ [\partial_N u] := \partial_N u^+ - \partial_N u^- = 0 & \text{on } S \end{cases} \quad (3.1)$$

is given by

$$u(x) = -b \int_S \partial_N \Gamma(x' - x) d\mathcal{H}^2(x'), \quad (3.2)$$

for $x \in \mathbb{R}^3 \setminus S$, where Γ is the fundamental solution in \mathbb{R}^3 of $\Delta \Gamma = \delta_0$. Moreover, such a solution is unique up to a harmonic map in \mathbb{R}^3 .

Here $\partial_N \Gamma = (\nabla_x \Gamma)N$. By (3.2), it holds that

$$\partial_i u(x) = -b \int_S \left(\frac{N_i}{|x - x'|^3} - 3 \frac{N \cdot (x - x')(x_i - x'_i)}{|x - x'|^5} \right) d\mathcal{H}^2(x'). \quad (3.3)$$

Lemma 3.2 is a particular case of its general vector counterpart, whose proof can be found in [39]. Its statement is given in the following lemma, where \mathbb{A} is a fourth-order elasticity tensor. So, the proof of Lemma 3.2 is straightforwardly obtained by letting $\mathbb{A} = \mathbb{I}_2$, the identity matrix.

LEMMA 3.3. *Let L be a Lipschitz closed curve in \mathbb{R}^3 and S a bounded Lipschitz surface with boundary L and unit normal N , as in assumption 3.1. Let $B \in \mathbb{R}^3$. The solution of*

$$\begin{cases} \operatorname{div} (\mathbb{A}\nabla w) = 0 & \text{in } \mathbb{R}^3 \setminus S \\ [w] := w^+ - w^- = B & \text{on } S \\ [(\mathbb{A}\nabla w)N] := ((\mathbb{A}\nabla w)N)^+ - ((\mathbb{A}\nabla w)N)^- = 0 & \text{on } S \end{cases} \quad (3.4)$$

is given componentwise by

$$w_i(x) = -B_j \int_S (\mathbb{A}\nabla\Gamma(x' - x) \cdot N(x'))_{ij} d\mathcal{H}^2(x'), \quad (3.5)$$

for $x \in \mathbb{R}^3 \setminus S$, where Γ is the solution in \mathbb{R}^3 of $\operatorname{div} (\mathbb{A}\nabla\Gamma) = \delta_0\mathbb{I}$.

3.1. Pointwise properties of gradients of harmonic maps

THEOREM 3.4 (Growth condition for the gradient). *Let L and S be as in lemma 3.2 and u be the explicit solution of (3.1) given by (3.2). If L is smooth, there exists a constant $c > 0$ depending on the curvature of L , such that it holds*

$$|\partial_i u(x)| \leq c|b|l \left(1 + \frac{1}{d(x, L)} \right). \quad (3.6)$$

Proof. Step 1. Let us first prove that the value of the derivative $\partial_i u(x)$ does not depend on the surface S appearing in (3.1). Let indeed S' be another smooth surface that does not contain the point x and has L as boundary. For simplicity, let us suppose it is disjoint from S . Let u' be the solution of (3.1) with S' replacing S and let A be the open set enclosed by S and S' . By formula (3.2), $(u - u')(x) = c + b \int_{\partial A} \partial'_N \Gamma(x' - x) d\mathcal{H}^2(x') = c + b\chi_A(x)$, the second equality being a consequence of the Divergence theorem. In particular, we see that $u - u'$ is constant in a neighbourhood of x , so that $\partial_i u(x) = \partial_i u'(x)$. By approximation, we can also extend this to the case of Lipschitz surface S' , and then to every rectifiable current S' with $\partial S' = L$ whose support is outside a neighbourhood of x .

Step 2. Let $d = d(x, L)$ be the distance from x to L , let κ be the maximum curvature of L , and let us denote by $R := \kappa^{-1}$ the minimal curvature radius. Let $B_R(x)$ be a ball with radius R and centre x , let P be the point in L such that $d = d(x, P)$, let O be the point on the line Px , on the x side, such that $d(O, P) = R$, and let $B_R(O)$ be a ball with radius R and centre O (see figure 1). Let $\pi_R : \mathbb{R}^3 \rightarrow \partial B_R(x)$ be the orthogonal projection onto the sphere $\partial B_R(x)$ and let L_R be the image of L by π_R . Let us consider the Lipschitz homotopy $\Phi : [0, 1] \times [0, l] \rightarrow \Omega$ such that $\Phi(0, [0, l]) = L$, $\Phi(1, [0, l]) = L_R$, and $\Phi(\cdot, t)$ is affine for all $t \in [0, l]$. Then $E := \Phi_{\#}[[0, 1] \times [0, l]]$ is a rectifiable current with boundary $L \cup L_R$. Let D be an integral current in $\partial B_R(x)$ with boundary $-L_R$. The rectifiable current $S' := E + D$ has boundary L , so we can consider the map u' solution of (3.1) with S replaced by S' .

Step 3. We claim that we can choose D in such a way that its total mass $M(D)$ is bounded by $3lR$, with l the total length of the curve L . Since L_R is an integral closed




Figure 1. Geometry

1-current, it decomposes as $L_R = \sum_i L_R^i$, with $\sum_i l_i = M(L_R) \leq l + 2\pi Rn \leq 3l$, where l_i is the length of L_R^i and n is the number of times that L enters inside B_R (in particular $n \leq \frac{l}{\pi R}$). Notice that the presence of the term $2\pi R$ is justified as follows: π_R is 1-Lipschitz outside $B_R(x)$ (which justifies the presence of l in the previous inequality), whereas we must estimate the inner part of L separately. Since the maximum curvature of L is assigned, it is seen that the part of L inside $B_R(x)$ is bounded and its projection on $B_R(x)$ does not exceed $2\pi R$. Having said that, by the isoperimetric inequality on $\partial B_R(x)$, it follows that

$$l_i^2 \geq 4\pi A_i - \frac{A_i^2}{R^2}, \tag{3.7}$$

with A_i the minimal area enclosed by L_R^i (see remark 3.5), so that $A_i \leq 2\pi R^2$. It follows from (3.7), by simple computations that $A_i \leq 2\pi R^2 - R\sqrt{4\pi^2 R^2 - l_i^2}$ and since $\sqrt{4\pi^2 R^2 - l_i^2} \geq 2\pi R - l_i$ if $l_i \leq 2\pi R$, one deduces by mere substitution that $A_i \leq Rl_i \wedge 2\pi R^2$. As a consequence, $\sum_i A_i \leq 3lR$, and hence $M(D) < 3lR$.

Step 4. Now, to compute $|\partial_i u(x)|$, we will use formula (3.3), integrating over $E + D$. Integration over D can be estimated as follows:

$$\begin{aligned} |\partial_i u(x)| &= \left| b \int_D \left(\frac{N_i}{|x - x'|^3} - 3 \frac{N \cdot (x - x')(x_i - x'_i)}{|x - x'|^5} \right) d\mathcal{H}^2(x') \right| \\ &\leq \frac{4|b|}{R^3} \int_D d\mathcal{H}^2(x') \leq \frac{12|b|l}{R^2}, \end{aligned} \tag{3.8}$$

since by Step 3, $M(D) \leq 3lR$. Let us now compute the integration over E . Observe that, by construction of E , it holds that its normal N at a point x' satisfies $N \perp (x - x')$, and hence by (3.3), we reduce to estimate

$$|b| \left| \int_E \frac{N_i}{|x - x'|^3} d\mathcal{H}^2(x') \right|.$$

Let E^+ (resp., E^-) be the part of E outside $B_R(x)$ (inside, respectively). The integration can be performed by polar coordinates (s, ρ) centred at x , where s can be seen as an arc parameter on L_R . The integral of (3.3) over E^+ is bounded by

$$\int_{L_R} ds \int_R^{+\infty} \frac{|b|}{\rho^3} d\rho \leq \frac{3l|b|}{2R^2}, \tag{3.9}$$

More delicate is the computation of the integral over E^- (see figure 1). Let L^- be the part of L inside $B_R(x)$. Let us assume that L^- has n connected components. Let L_1^- be one of this components. By the regularity hypotheses on L and since $B_R(O)$ is tangent to L at P , one has that L stays outside $B_R(O)$. Let P_1 and P_2 be the two points on $\partial B_R(O) \cap \partial B_R(x)$ belonging to the plane containing O, x , and the tangent vector τ to L at P . Due to the boundedness of the curvature of L , it can be easily observed that the integral of (3.3) on the component L_1^- of L^- is maximal when L_1^- is planar and remains in the circumference obtained intersecting $B_R(O)$ with the plane containing O, x , and the tangent vector τ (the sheet plane in figure 1). In such a case L_1^- is the arc of circumference passing through P and with endpoints P_1 and P_2 . Moreover, E^- is planar and is contained into two half-planes Π and Π' passing through P , containing the tangent τ , and with boundaries the lines $\overline{P_1P}$ and $\overline{P_2P}$, respectively (in figure 1).

Now we estimate the integral on every connected components of L^- , and then multiply the obtained value by n in order to get a right estimate. Since the half-planes Π and Π' are constructed in such a way that E^- is contained in their union, the integral

$$\int_{E^-} \frac{|b|}{d(x, x')^3} d\mathcal{H}^2(x'),$$

is bounded by the same integral over $\Pi \cup \Pi'$. In particular, by symmetry, it holds

$$\int_{E^-} \frac{|b|}{d(x, x')^3} d\mathcal{H}^2(x') \leq 2 \int_{\Pi} \frac{|b|}{d(x, x')^3} d\mathcal{H}^2(x').$$

If the distance $d(x, \Pi) = \delta$ the right-hand side is easily computed by integrating in planar polar coordinates with $\delta \leq r = d(x, x') < \infty$, and is equal to $2\pi|b|/\delta$. Since by construction $\delta \geq 2^{-1/2}d(x, L)$, it holds

$$\int_{E^-} \frac{|b|}{d(x, x')^3} d\mathcal{H}^2(x') \leq |b| \frac{4\pi\sqrt{2}}{d(x, L)}.$$

Now we have to multiply this by n , the number of connected components of L^- . We claim that $n \leq l/(\pi R)$. Indeed, consider a single connected component and one of its

point P inside $B_R(x)$. With elementary geometric arguments, it is seen that due to the boundedness of the curvature, as observed, it could stay inside $B_R(x) \setminus B_R(O)$ for a length at most πR . On the other hand, if it goes out of $B_R(x) \setminus B_R(O)$, then it cannot come back before spending a length of at least πR outside. Then the claim follows, and

$$\int_{E^-} \frac{|b|}{d(x, x')^3} d\mathcal{H}^2(x') \leq \frac{4\sqrt{2}}{R} \frac{|b|l}{d(x, L)}.$$

Summing all the bounds obtained so far, we finally get

$$|\partial_i u(x)| \leq 4\sqrt{2} \frac{|b|l\kappa}{d(x, L)} + 12|b|l\kappa^2 + \frac{3}{2}l|b|\kappa^2, \tag{3.10}$$

from which the thesis follows. □

REMARK 3.5. In the previous proof, we can choose the sets A_i with minimal area as follows. Let L_R^i be a simple loop in $B_R(x)$, let $P \notin L_R^i$ be a point on $\partial B_R(x)$. We construct a homotopy $\Psi^P : [0, 1] \times [0, 2\pi] \rightarrow \partial B_R(x)$ that satisfies $\Psi^P(0, \cdot) \equiv P$ and $\Psi^P(1, [0, 2\pi]) = L_R^i$, and we can consider the current $\Psi_\#^P[[0, 1] \times [0, 2\pi]]$. Then we can set $A^i := \Psi_\#^P[[0, 1] \times [0, 2\pi]]$, where P is chosen in such a way that $\Psi_\#^P[[0, 1] \times [0, 2\pi]]$ has minimal mass.

REMARK 3.6. In theorem 3.4, we also proved that the integral in (3.2) does not depend on the particular surface S , but only on its boundary L .

LEMMA 3.7. *Let $b \in 2\pi\mathbb{Z}$. Then the solution u of (3.1) in (3.2) belongs to $C^\infty(\Omega \setminus L, \mathbb{T})$ and it is harmonic in $\Omega \setminus L$.*

Proof. As we have proved in Step 1 of theorem 3.4, if we choose a surface S' with boundary L disjoint from S , and denote by u' the corresponding solution of (3.1), then $u - u' = b\chi_A$, with A the open set with boundary $S \cup S'$. Since $b \in 2\pi\mathbb{Z}$ we see that $u = u'$ as a map into \mathbb{T} . Moreover, if $x \notin S$ then u is smooth at x , so in particular, up to change the surface S , we obtain that it belongs to $C^\infty(\Omega \setminus L, \mathbb{T})$ and u is harmonic at x for all $x \notin L$. □

COROLLARY 3.8. *Let L be the union of $N > 0$ smooth closed curves L_k , let S be the union of the corresponding surfaces S_k with boundary L_k respectively, and let u be the solution to (3.1) given by (3.2). Then (3.6) holds true.*

Proof. Actually, the same proof as of theorem 3.4 applies. □

COROLLARY 3.9. *Let L and S be as in lemma 3.3 and w be the explicit solution of (3.4). If L is smooth, there exists a constant $c > 0$ depending on the curvature of L , such that it holds*

$$|\partial_j w_i(x)| \leq c|B|l \left(1 + \frac{1}{d(x, L)} \right), \quad 1 \leq i, j \leq 3. \tag{3.11}$$

THEOREM 3.10 (Finer description of the singularity). *Let S and L be as in assumption 3.1. Let u be the solution to*

$$\begin{cases} \Delta u = 0 & \text{in } \mathbb{R}^3 \setminus S \\ u^+ - u^- = 1 & \text{on } S \\ \partial_N u^+ - \partial_N u^- = 0 & \text{on } S, \end{cases} \tag{3.12}$$

given by (3.2) with $b = 1$. Then, if U is a tubular neighbourhood of L , for all $(\rho, \theta, \tau) \in U$ with $\theta \neq 0$, it holds

- (i) *there exists the limit $\lim_{\epsilon \rightarrow 0^+} u(\epsilon\rho, \theta, \tau) = u(0^+, \theta, \tau) = \frac{2\theta}{2\pi} + c$, where c is a constant independent of θ .*
- (ii) *$\lim_{\epsilon \rightarrow 0^+} |\partial_\tau u(\epsilon\rho, \theta, \tau)| < c < +\infty$ for some constant $c > 0$ that depends only on the curve L .*
- (iii) *$\lim_{\epsilon \rightarrow 0^+} |\partial_\rho u(\epsilon\rho, \theta, \tau)| < c < +\infty$ for some constant $c > 0$ that depends only on the curve L .*

Proof. With no loss of generality, we suppose that the curve L which represents the boundary of S passes through the origin of a Euclidean coordinate system where it is tangent to the z -axis. Moreover, we choose the coordinates x_1 and x_2 in such a way that $x_1 = \rho \cos \theta$ and $x_2 = \rho \sin \theta$, so it follows that the point $(\epsilon\rho, \theta, z)$ coincides with $(\epsilon x_1, \epsilon x_2, z)$. For simplicity, we take $z = 0$ and denote $x = (x_1, x_2, 0)$, while S is orthogonal to the x_2 -axis at 0. From lemma 3.2, we have the following explicit formula

$$u(\epsilon\rho, \theta, 0) = u(\epsilon x_1, \epsilon x_2, 0) = - \int_S \partial_N \Gamma(x' - \epsilon x, y' - \epsilon y, z') d\mathcal{H}^2(x', y', z').$$

By the change of variables $(\epsilon x''_1, \epsilon x''_2, \epsilon z'') = (x'_1, x'_2, z')$ we obtain

$$u(\epsilon x_1, \epsilon x_2, 0) = - \int_{1/\epsilon S} \partial_N \Gamma(x''_1 - x_1, x''_2 - x_2, z'') d\mathcal{H}^2(x''_1, x''_2, z''),$$

where we have used the explicit expression of Γ . Letting ϵ go to zero, we obtain

$$\lim_{\epsilon \rightarrow 0^+} u(\epsilon\rho, \theta, z) = - \int_{\Pi_0} \partial_N \Gamma(x'' - x) d\mathcal{H}^2(x''),$$

where Π_0 is the half-plane $\{x_1 > 0, x_2 = 0\}$ and we have used the shorter notation $x'' = (x''_1, x''_2, z'')$. Thanks to lemma 3.2, we see that the right-hand side coincides with $\hat{u}(\rho, \theta, z)$, where \hat{u} is the solution of (3.12) with $S = \Pi_0$. But it is well known that such solution is given by, in cylindrical coordinates, $\hat{u}(\rho, \theta, z) = \frac{\theta}{2\pi} + c$ for a fixed constant c . In particular we have $\lim_{\epsilon \rightarrow 0^+} u(\epsilon\rho, \theta, z) = \frac{\theta}{2\pi} + c$.

To prove statement (ii), we use the explicit expression (3.3), which reads, after the change of variables $x' = \epsilon x'$ (here, again with abuse of notations, $\partial_z u = \partial_\tau u$ at

$x = 0$),

$$\partial_z u(\epsilon x_1, \epsilon x_2, 0) = -\frac{1}{\epsilon} \int_{1/\epsilon S} \left(\frac{N_z}{|x - x''|^3} - 3 \frac{N \cdot (x - x'')(z - z'')}{|x - x''|^5} \right) d\mathcal{H}^2(x''). \tag{3.13}$$

We fix $\delta > 0$ and consider the ball B_ϵ with centre $(\epsilon x_1, \epsilon x_2, 0)$ and radius δ . We then write the last integral as

$$\begin{aligned} & -\frac{1}{\epsilon} \int_{1/\epsilon S \cap B_\epsilon} \left(\frac{N_z}{|x - x''|^3} - 3 \frac{N \cdot (x - x'')(z - z'')}{|x - x''|^5} \right) d\mathcal{H}^2(x'') \\ & -\frac{1}{\epsilon} \int_{1/\epsilon S \cap B_\epsilon^c} \left(\frac{N_z}{|x - x''|^3} - 3 \frac{N \cdot (x - x'')(z - z'')}{|x - x''|^5} \right) d\mathcal{H}^2(x''), \end{aligned}$$

and thanks to remark 3.6, up to choosing δ small enough, we can assume that the normal N to S at x' is orthogonal to the vector $(\epsilon x - x')$ for all $x' \in B_\epsilon$, that is, to $(x - x'')$ in $1/\epsilon B_\epsilon$, so that the integral above becomes

$$\begin{aligned} & -\frac{1}{\epsilon} \int_{1/\epsilon S \cap B_\epsilon} \frac{N_z}{|x - x''|^3} d\mathcal{H}^2(x'') \\ & -\frac{1}{\epsilon} \int_{1/\epsilon S \cap B_\epsilon^c} \left(\frac{N_z}{|x - x''|^3} - 3 \frac{N \cdot (x - x'')(z - z'')}{|x - x''|^5} \right) d\mathcal{H}^2(x''). \end{aligned} \tag{3.14}$$

Let us now estimate the second term in (3.14). In B_ϵ^c it holds $|\epsilon x - x'| > \delta$, that is, $|x - x''| > \epsilon^{-1}\delta$, so it is easy to see that this term can be estimated by

$$\frac{\mathcal{H}^2(S)}{\delta^3} \leq (\gamma + 1) \frac{l^2}{\delta^3},$$

where l is the length of L and $\gamma > 0$ is the constant of the isoperimetric inequality, since S can be chosen arbitrarily.

It remains to estimate the first term in (3.14). Let us consider the plane Π passing through 0 and tangent to the vector \bar{z} and to $x - \bar{0}$ (the unit vector tangent to the z -axis and the vector passing through x and 0, respectively). Let Π^+ be the half-plane in Π bounded by the axis \hat{z} and not containing the point x . Thanks to the smoothness of L , we can assume that there exists a smooth one-to-one map $\Phi : \Pi^+ \cap B_\epsilon \rightarrow S \cap B_\epsilon$, so that also $N \circ \Phi : \Pi^+ \ni \hat{x}' \mapsto N(x')$ is smooth, and then in $B_\epsilon \cap \Pi^+$ we can use the Taylor expansion of $N \circ \Phi$ at 0. Going back to the variable $x' = \epsilon x''$ (and $\hat{x}' := \epsilon \hat{x}''$), we find that the first term in (3.14) reads

$$-\int_{S \cap B_\epsilon} \frac{N(x')}{|\epsilon x - x'|^3} d\mathcal{H}^2(x') = -\int_{S \cap B_\epsilon} \frac{\nabla^2 N(0) \hat{x}' \cdot \hat{x}'}{|\epsilon x - x'|^3} + \frac{r_N(|\hat{x}'|^2)}{|\epsilon x - x'|^3} d\mathcal{H}^2(x').$$

The Taylor expansion of Φ at 0 provides $x' = \hat{x}' + \nabla^2 \Phi(0) \hat{x}' \cdot \hat{x}' + r_\Phi(\hat{x}')$ and if δ is small enough, we can assume that $|\nabla^2 \Phi(0) \hat{x}' \cdot \hat{x}' + r_\Phi(\hat{x}')| < 1/2|\hat{x}'|$. Note that, since L is smooth, we can find such a $\delta > 0$ satisfying the last inequality globally, that is, δ is independent of the point x . In particular, we find $|\epsilon x - x'| > |\epsilon x - \hat{x}'| -$

$|\nabla^2\Phi(0)\hat{x}' \cdot \hat{x}' + r_\Phi(\hat{x}')| > |\hat{x}'| - 1/2|\hat{x}'| = 1/2|\hat{x}'|$ for all $\epsilon > 0$, so that the integral is bounded by

$$\int_{\Pi^+ \cap B_\epsilon} \frac{|\nabla^2 N(0)\hat{x}' \cdot \hat{x}'|}{|\hat{x}'|^3} + \frac{r_N(|\hat{x}'|^2)}{|\hat{x}'|^3} d\mathcal{H}^2(x'), \tag{3.15}$$

and taking into account that $\delta > 0$ can be small as we want, we can also assume that $|r_N(\hat{x}')| < |\nabla^2 N(0)\hat{x}' \cdot \hat{x}'|$, whereby the last integral can be estimated by

$$C \int_{\Pi^+ \cap B_\epsilon} \frac{1}{|\hat{x}'|} d\mathcal{H}^2(\hat{x}'),$$

where the constant C is independent of δ and x . Taking the limit as $\epsilon \rightarrow 0$, we infer, by the monotone convergence theorem,

$$C \int_{\Pi^+ \cap B(0,\delta)} \frac{1}{|\hat{x}'|} d\mathcal{H}^2(\hat{x}'),$$

which is uniformly bounded. Now, since the value of δ is independent of the point x but only depends on the geometry of the curve L , we achieved the proof.

Statement (iii) can be proved taking into account that choosing δ small enough a formula similar to (3.14) holds, and then arguing as for statement (ii). \square

REMARK 3.11. Let us point out that theorem 3.10 still holds true if we do not assume that L is connected. Indeed if L is the union of a finite family of smooth closed curves, the surface S will be the union of a finite family of smooth surfaces and the arguments used in the proof of theorem 3.10 still work.

REMARK 3.12. The curve regularity required in theorem 3.10 is $W^{3,\infty}$, because of estimates such as (3.15). Note also that the constant L depends on the curve curvature.

3.2. Distributional properties of gradients of harmonic maps

In this section, and in the following, we often deal with BV-functions u (functions of bounded variation). In this case, if not otherwise advised, we will always denote by Du the distributional derivative of u , and by ∇u the absolutely continuous part of Du with respect to the Lebesgue measure.

THEOREM 3.13. *Let L be a closed Lipschitz curve in Ω and let $b \in 2\pi\mathbb{Z}^3$. Then for any Lipschitz surface S with boundary L as in assumption 3.1, every solution u to (3.1) belongs to $BV^p(\Omega, \mathbb{R}^3)$ with $1 \leq p < 3/2$, satisfies $\text{Div } \nabla u = 0$ and $-\text{Curl } \nabla u = b \otimes \mathcal{L}$ as distributions, with ∇u the part of the gradient of u that is absolutely continuous with respect to the Lebesgue measure.*

Proof. Let u be a solution to (3.1) and let us first assume L smooth. By (3.6), $\nabla u \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ for $p < 2$ (see the second statement of theorem 6.3). It has been shown that u is smooth outside S where it has a jump of amplitude b . In particular, u belongs to $SBV(\Omega, \mathbb{R}^3)$ and its distributional derivative is given by

$$\langle Du, \varphi \rangle := -\langle u, \text{div } \varphi \rangle = S(\varphi) + \langle \nabla u, \varphi \rangle, \tag{3.16}$$

for all $\varphi \in C_c^\infty(\Omega, \mathbb{R}^{3 \times 3})$, where S denotes the distribution $S(\varphi) = - \int_S N_j b_i \varphi_{ij} d\mathcal{H}^2$.

Let us prove that $-\text{Curl } \nabla u = b \otimes \mathcal{L}$. To this aim, let us take $\psi \in \mathcal{C}_c^\infty(\Omega, \mathbb{R}^{3 \times 3})$ and write

$$\begin{aligned}
 -\langle \text{Curl } \nabla u, \psi \rangle &:= -\langle \nabla u, \text{Curl } \psi \rangle = -\langle Du, \text{Curl } \psi \rangle + S(\text{Curl } \psi) \\
 &= \int_L \tau_j b_i \psi_{ij} d\mathcal{H}^1 = b \otimes \mathcal{L}(\psi),
 \end{aligned} \tag{3.17}$$

where the second equality follows from (3.16) with $\varphi = \text{Curl } \psi$, and the third one by Stokes theorem. We now prove that $\text{Div } \nabla u = 0$. We take $\psi \in \mathcal{C}_c^\infty(\Omega, \mathbb{R}^3)$ and write

$$-\langle \text{Div } \nabla u, \psi \rangle := \langle \nabla u, \nabla \psi \rangle = \langle Du, \nabla \psi \rangle - S(\nabla \psi), \tag{3.18}$$

and using the explicit formula (3.2) for u , we obtain

$$\begin{aligned}
 \langle Du, \nabla \psi \rangle &= b_k \left\langle D_i \int_S \partial_N \Gamma_k(x' - \cdot) d\mathcal{H}^2(x'), D_i \psi \right\rangle \\
 &= - \int_S b_k \langle \Delta \Gamma_k(x' - \cdot), D_j \psi N_j \rangle d\mathcal{H}^2(x') \\
 &= -b_k \int_S \partial_N \psi_k(x') d\mathcal{H}^2(x') = S(\nabla \psi),
 \end{aligned}$$

so that plugging the last identity in (3.18), we obtain $\text{Div } \nabla u = 0$.

Let us now treat the general case. We proceed by approximation, so let L_n be a sequence of curves converging uniformly and in the sense of currents to L , let S_n be surfaces converging in the same sense to S , and let u_n be the corresponding solutions. Let V_n be the solution to the system

$$\begin{cases} -\text{Curl } U &= -b \otimes \mathcal{L}_n & \text{in } \Omega \\ \text{div } U &= 0 & \text{in } \Omega \\ UN &= 0 & \text{on } \partial\Omega, \end{cases} \tag{3.19}$$

which, by theorem 6.3 below also satisfies $\|U\|_{L^p} \leq c|\mu|(\Omega)$, with $c = c(\Omega)$, so that we infer $\|V_n\|_{L^p(\Omega)} \leq |b| |\mathcal{L}_n|$ for $1 \leq p < 3/2$. So far, we have proved that ∇u_n differs from $-V_n$ by the gradient of a harmonic map whose boundary datum at $\partial\Omega$ are bounded since $d(L_n, \partial\Omega) > \delta > 0$ (this can be seen computing $\nabla u_n N$ by formula (3.3)). In particular, we find that there are constants $c_1, c_2 > 0$ such that $\|\nabla u_n\|_{L^p(\Omega)} \leq c_1 |b| |\mathcal{L}_n| + c_2$ for $1 \leq p < 3/2$. Now it is easy to see that $u_n \rightharpoonup u$ weakly* in $BV^p(\Omega, \mathbb{R}^3)$, and the conclusion easily follows. \square

REMARK 3.14. In order to prove that $\text{Div } \nabla u = 0$, we might also argue as follows. Let $\hat{S} \supset S$ such that \hat{S} separates Ω in two parts Ω^- and Ω^+ . Then for every test function $\varphi \in \mathcal{C}_c^\infty(\Omega, \mathbb{R}^3)$ it holds

$$\begin{aligned}
 \int_\Omega \nabla u \nabla \varphi dx &= \int_{\Omega^+} \nabla u \nabla \varphi dx + \int_{\Omega^-} \nabla u \nabla \varphi dx = \\
 &- \int_{\Omega^+} \text{Div } \nabla u \varphi dx - \int_{\Omega^-} \text{Div } \nabla u \varphi dx + \int_{\hat{S}^+} \partial_N u^+ \varphi dx - \int_{\hat{S}^-} \partial_N u^- \varphi dx = 0.
 \end{aligned}$$

Moreover, if the curve L is smooth, it can be proved that $\nabla u \in L^p(\Omega, \mathbb{R}^{3 \times 3})$ for all $1 \leq p < 2$. This will be addressed in theorem 6.3 below.

REMARK 3.15. The statement of theorem 3.13 readily applies to the case of L being a finite union of Lipschitz curves.

According to corollary 3.3, the counterpart of theorem 3.13 for a vector-valued displacement in linear elasticity reads as follows:

COROLLARY 3.16. *Let $L \subset \Omega$ be the union of a finite number of smooth dislocation loops and $S \subset \Omega$ a smooth surface enclosed by L . Referring to lemma 3.3, let w be the solution of*

$$-\operatorname{div}(\mathbb{A}\nabla w) = 0 \quad \text{in } \mathbb{R}^3 \setminus S, \quad [w] = B \in 2\pi\mathbb{Z}^3, \quad [(\mathbb{A}\nabla w)N] = 0 \quad \text{on } S,$$

given by (3.5). Then $w \in SBV(\Omega, \mathbb{R}^3)$, $\nabla w \in L^p(\Omega, \mathbb{R}^3)$ for $1 \leq p < 2$ and $-\operatorname{Curl} \nabla w = B \otimes \mathcal{L}$ in the distributional sense, where ∇w is the absolutely continuous part of the distributional derivative Dw in Ω . Moreover, $-\operatorname{div}(\mathbb{A}\nabla w) = 0$ in $\mathbb{R}^3 \setminus C$, $w \in C^\infty(\Omega \setminus \mathcal{L}, \mathbb{R}^3)$ and it holds

$$|\nabla w(x)| \leq c|B|l \left(1 + \frac{1}{d(x, L)}\right), \tag{3.20}$$

with c a constant depending on the maximal curvature of L , and l its length.

The bound (3.20) is proved as in theorem 3.4, since \mathbb{A} is a bounded tensor.

4. Main result: boundary of the graph of \mathbb{T}^3 -valued harmonic maps

Preliminary notions. We introduce the following notation. For all $b \in \mathbb{R}^3$, we define the 1-current $\vec{b} \in \mathcal{D}_1(\mathbb{T}^3)$ as

$$\vec{b}(\omega) := -\frac{1}{2\pi} \int_0^{2\pi} \left\langle \omega \left(\frac{b_1\theta}{2\pi}, \frac{b_2\theta}{2\pi}, \frac{b_3\theta}{2\pi} \right), b \right\rangle d\theta, \tag{4.1}$$

for any 1-form $\omega \in \mathcal{D}^1(\mathbb{T}^3)$. It is easy to see that $M(\vec{b}) = |b|$. The fact that we are on the torus, that is, ω is 2π -periodic on \mathbb{R}^3 , implies that \vec{b} is a closed current whenever $b \in 2\pi\mathbb{Z}^3$. Moreover, it is convenient to define, for all $b \in \mathbb{R}^3$ and all $r \in \mathbb{R}^3$, the 1-current $\vec{b}_r \in \mathcal{D}_1(\mathbb{T}^3)$ as

$$\vec{b}_r(\omega) := -\frac{1}{2\pi} \int_0^{2\pi} \left\langle \omega \left(r_1 + \frac{b_1\theta}{2\pi}, r_2 + \frac{b_2\theta}{2\pi}, r_3 + \frac{b_3\theta}{2\pi} \right), b \right\rangle d\theta, \tag{4.2}$$

for any 1-form $\omega \in \mathcal{D}^1(\mathbb{T}^3)$. Also in this case $M(\vec{b}_r) = |b|$ for all $r \in \mathbb{R}^3$. Note that if $b \in 2\pi\mathbb{Z}^3$ and there is a real number δ such that $b = \delta r$, then the currents $\vec{b} = \vec{b}_r$ thanks to the periodicity of the forms in $\mathcal{D}^1(\mathbb{T}^3)$.

We denote by $\mathcal{L} \wedge \vec{b}$ the 2-current in $\Omega \times \mathbb{T}^3$ defined as

$$\mathcal{L} \wedge \vec{b}(\omega) = -\frac{1}{2\pi} \int_L \int_0^{2\pi} \left\langle \omega \left(x, \frac{b\theta}{2\pi} \right), \bar{\tau} \wedge \vec{b} \right\rangle d\theta d\mathcal{H}^1(x), \tag{4.3}$$

for any 2-form $\omega \in \mathcal{D}^2(\Omega \times \mathbb{T}^3)$ (we mean $\bar{\tau} = (\vec{\tau}, 0) \in \mathbb{R}^3 \times \mathbb{R}^3$, the tangent vector to L in $\Omega \times \mathbb{R}^3$, and $\vec{b} = (0, b) \in \mathbb{R}^3 \times \mathbb{R}^3$, with $\vec{\tau}$ the tangent vector to L in Ω).

The cylindrical neighbourhood. Let L be a closed loop of class \mathcal{C}^2 . There is a cylindrical neighbourhood D_R of L , with cylindrical coordinates $(\rho, \theta, z) \in [0, R] \times [0, 2\pi] \times [0, l] / \sim$, where \sim means that the coordinate $\theta = 0$ (and $z = 0$) is identified with $\theta = 2\pi$ (resp. $z = l$). The neighbourhood D_R is also parametrized by the coordinates (x, y, z) by setting $x = \rho \cos \theta$ and $y = \rho \sin \theta$. Let S be a smooth surface with boundary L and such that $S \cap D_R$ coincides with the set $\{\theta = 0\}$. We suppose assumption 3.1 on the regularity of S . The tangent space to ∂D_R is a 2-dimensional plane with orthonormal basis $\{\sigma, \tau\}$, with τ representing the vector tangent to the curve, and then corresponding to the z coordinate, and σ tangent to the section (orthogonal to L) of ∂D_R , namely corresponding to the coordinate θ .

4.1. Statement of the main theorem

We are ready to prove the following:

THEOREM 4.1. *Let S be a smooth surface in Ω whose boundary L is a smooth and closed curve in Ω , both satisfying assumption 3.1. Let $b = (b_1, b_2, b_3) \in 2\pi\mathbb{Z}^3$ and let $u = (u_1, u_2, u_3) : \Omega \rightarrow \mathbb{R}^3$ be the map where u_i is given by (3.2) with $b = b_i$. Then \mathcal{G}_u is an integral current in $\mathcal{D}_3(\Omega \times \mathbb{T}^3)$ and its boundary is given by*

$$\partial \mathcal{G}_u(\omega) = \mathcal{L} \wedge \vec{b}(\omega), \tag{4.4}$$

for all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{T}^3)$.

Let $\Phi_\theta(x) := (x, b_1\theta/2\pi, b_2\theta/2\pi, b_3\theta/2\pi)$ and define the tensor test function φ^ω associated with any form $\omega \in \mathcal{D}^2(\Omega \times \mathbb{T}^3)$ componentwise as

$$\varphi_{\alpha\beta}^\omega := \int_0^{2\pi} \frac{\langle \omega \circ \Phi_\theta, e_\alpha \wedge e_\beta \rangle}{2\pi} d\theta, \tag{4.5}$$

where $\alpha, \beta = 1, 2, 3$. Then, our main result can be restated as

$$\partial \mathcal{G}_u(\omega) = \Lambda_{\mathcal{L}}(\varphi^\omega) := \mathcal{L} \otimes b(\varphi^\omega), \tag{4.6}$$

for all $\omega \in \mathcal{D}^3(\Omega \times \mathbb{T}^3)$, where φ^ω is defined by (4.5).

REMARK 4.2. For all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{T}^3)$ let us write $\omega = \sum_{i=0}^2 \omega^i$ where we have set

$$\omega^i := \sum_{\alpha:|\alpha|=i} \omega_{\alpha\beta} dx^\alpha \wedge dy^\beta. \tag{4.7}$$

In other words, ω^i represents the components of ω whose coefficient $dx^\alpha \wedge dy^\beta$ has a i -dimensional horizontal component (dx^α) and a $2 - i$ dimensional vertical

component (dy^β) . It is seen that $\partial\mathcal{G}_u$ only depends on the mixed components of ω , that is, on the component ω^1 in (4.7).

COROLLARY 4.3. *Let S be a Lipschitz surface in Ω whose boundary L is a Lipschitz and closed curve in Ω . Let $b = (b_1, b_2, b_3) \in 2\pi\mathbb{Z}^3$ and let $u = (u_1, u_2, u_3) : \Omega \rightarrow \mathbb{R}^3$ be a map with u_i satisfying (3.2) with $b = b_i$. Then \mathcal{G}_u is an integral current in $\mathcal{D}_3(\Omega \times \mathbb{T}^3)$ and (4.4) holds.*

Proof. We proceed by approximation. Let $\{L_k\}_{k>0}$ be a sequence of smooth closed curves approximating L (uniformly and in the sense of 1-currents) and let $\{S_k\}_{k>0}$ be smooth surfaces with boundary $\{L_k\}_{k>0}$ and converging (uniformly and in the sense of currents) to S . Let u_k be maps as in theorem 4.1 with L replaced by L_k and S replaced by S_k . Thanks to the uniform convergence of S_k to S and using formula (3.2) we see that u_k converges pointwise to u , and then strongly in $L^p(\Omega, \mathbb{T}^3)$. Since L_k are converging uniformly to L whose length is finite, the lengths of L_k are uniformly bounded so the same argument employed in theorem 3.13 gives a uniform bound in $L^p(\Omega)$, with $p < 3/2$, for the 1×1 minors of Du_k , while the higher-order minors are all null. Therefore, there are maps $v_\alpha^\beta \in L^p(\Omega)$ such that, up to a subsequence, $M_\alpha^\beta(Du_k) \rightharpoonup v_\alpha^\beta$ weakly in $L^p(\Omega)$. Finally, the lengths of L_k being uniformly bounded, theorem 4.1 provides a uniform bound on the masses of $\partial\mathcal{G}_{u_k}$. Now theorem 2.3 applies and implies that $u \in \mathcal{A}_p(\Omega, \mathbb{T}^3)$. In particular, we have that $u_k \rightharpoonup u$ weakly in $\mathcal{A}_p(\Omega, \mathbb{T}^3)$, thus lemma 2.2 implies that $\partial\mathcal{G}_{u_k} \rightharpoonup \partial\mathcal{G}_u$ as currents, and the fact that for u_k the explicit form (4.4) holds true implies that it holds also at the limit, concluding the proof. □

4.2. Proof of theorem 4.1

In the sequel, we will use the notation

$$\Phi := Id \times u : \Omega \rightarrow \Omega \times \mathbb{T}^3.$$

Before proving theorem 4.1, we state the following preliminary fact:

LEMMA 4.4. *Let u be as in theorem 4.1. Then $u \in \mathcal{A}_p(\Omega, \mathbb{T}^3)$ for all $1 \leq p < 2$.*

Proof. The fact that $u \in W^{1,p}(\Omega, \mathbb{T}^3)$ for all $1 \leq p < 2$ can be proved as in theorem 6.3 below. Moreover, lemma 3.7 shows that u is well-defined in \mathbb{T}^3 . In order to prove that it belongs to $\mathcal{A}_p(\Omega, \mathbb{T}^3)$, we need to show that all its minors $M_\alpha^\beta(Du)$ belong to $L^p(\Omega)$. Thanks to theorem 3.4 it is easy to see that every 1×1 -minor belongs to $L^p(\Omega)$. Moreover, from lemma 3.2 we have that u_1, u_2 , and u_3 differ by a multiplicative constant so that the rows of the matrix Du are linearly dependent. In particular, all the minors $>1 \times 1$ vanish, and the thesis follows. □

Proof of theorem 4.1. Let u_ϵ be the restriction of the map u to $\Omega_\epsilon := \Omega \setminus \bar{D}_\epsilon$, $u_\epsilon := u|_{\Omega_\epsilon}$, where $D_\epsilon := \{(\rho, \theta, z) \in [0, R] \times [0, 2\pi] \times [0, l] / \sim : \rho < \epsilon\}$ is a cylindrical neighbourhood of L . The graph \mathcal{G}_{u_ϵ} is the restriction of the graph \mathcal{G}_u to the open set $\Omega_\epsilon \times \mathbb{T}^3$. Formula (2.6) and the Dominated Convergence Theorem readily

implies that $\mathcal{G}_{u_\epsilon} \rightharpoonup \mathcal{G}_u$ as currents. As a consequence, we find

$$\partial \mathcal{G}_{u_\epsilon} \rightharpoonup \partial \mathcal{G}_u.$$

In order to compute explicitly the boundary of \mathcal{G}_u , we write $\partial \mathcal{G}_{u_\epsilon}(\omega) = \mathcal{G}_{u_\epsilon}(d\omega)$, for $\omega \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$. Lemma 3.7 implies that u is smooth outside a neighbourhood of L so that we can apply the Stokes Theorem and find

$$\partial \mathcal{G}_{u_\epsilon}(\omega) = \int_{\partial D_\epsilon} \left\langle \omega \circ \Phi, \frac{\partial \Phi}{\partial \sigma} \wedge \frac{\partial \Phi}{\partial \tau} \right\rangle d\mathcal{H}^2,$$

where (σ, τ) is an orthogonal coordinate system in the tangent space to ∂D_ϵ . The gradient of Φ reads

$$(D\Phi) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ \frac{\partial u_1}{\partial x_1} & \frac{\partial u_1}{\partial x_2} & \frac{\partial u_1}{\partial x_3} \\ \frac{\partial u_2}{\partial x_1} & \frac{\partial u_2}{\partial x_2} & \frac{\partial u_2}{\partial x_3} \\ \frac{\partial u_3}{\partial x_1} & \frac{\partial u_3}{\partial x_2} & \frac{\partial u_3}{\partial x_3} \end{pmatrix}. \tag{4.8}$$

Let $\partial D_\epsilon \cong [0, 2\pi] \times [0, l] / \sim$ for all $(\theta, \tau) \in \partial D_\epsilon$. In the coordinate system $(\rho, \sigma, \tau, y_1, y_2, y_3)$ it holds

$$(D(\Phi_{\perp \partial D_\epsilon})) = \left(\frac{\partial \Phi}{\partial \sigma}, \frac{\partial \Phi}{\partial \tau} \right) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \\ \frac{\partial u_1}{\partial \sigma} & \frac{\partial u_1}{\partial \tau} \\ \frac{\partial u_2}{\partial \sigma} & \frac{\partial u_2}{\partial \tau} \\ \frac{\partial u_3}{\partial \sigma} & \frac{\partial u_3}{\partial \tau} \end{pmatrix}. \tag{4.9}$$

If $\omega = \omega_{ij} dz^i \wedge dz^j$, with $1 \leq i < j \leq 6$, where we have defined $z_1 = \rho$, $z_2 = \sigma$, $z_3 = \tau$, and $z_{k+3} = y_k$ for $k = 1, 2, 3$, we can write

$$\int_{\partial D_\epsilon} \left\langle \omega \circ \Phi, \frac{\partial \Phi}{\partial \sigma} \wedge \frac{\partial \Phi}{\partial \tau} \right\rangle d\mathcal{H}^2(x) = \int_{\partial D_\epsilon} \omega_{ij}(x, u(x)) \tilde{M}_i^j(D(\Phi_{\perp \partial D_\epsilon}(x))) d\mathcal{H}^2(x), \tag{4.10}$$

with $\tilde{M}_i^j(D(\Phi_{\perp \partial D_\epsilon}(x)))$ being the minor of $D(\Phi_{\perp \partial D_\epsilon})$ given by the i -th and j -th rows. From (4.9), we see that the (2×2) -minors of $D(\Phi_{\perp \partial D_\epsilon})$ which are nonzero are the only ones involving either the second or third row. So (4.10) reads

$$\int_{\partial D_\epsilon} \left(\omega_{23}(x, u(x)) + \sum_{k=4}^6 \left(\omega_{2k}(x, u(x)) \frac{\partial u_{k-3}}{\partial \tau}(x) - \omega_{3k}(x, u(x)) \frac{\partial u_{k-3}}{\partial \sigma}(x) \right) \right) d\mathcal{H}^2(x)$$

$$\begin{aligned}
 &= \int_{\partial D_\epsilon} \omega_{23}(x, u(x)) d\mathcal{H}^2(x) + \sum_{k=4}^6 \int_0^{2\pi} \int_0^l \epsilon \tilde{\omega}_{2k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) \frac{\partial u_{k-3}}{\partial \tau}(\epsilon, \theta, \tau) d\tau d\theta \\
 &- \sum_{k=4}^6 \int_0^l \int_0^{2\pi} \epsilon \tilde{\omega}_{3k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) \frac{\partial u_{k-3}}{\partial \sigma}(\epsilon, \theta, \tau) d\theta d\tau, \tag{4.11}
 \end{aligned}$$

where $\tilde{\omega} := \omega \det \Psi$, with $\Psi : [0, \epsilon] \times [0, 2\pi] \times [0, l] \rightarrow D_\epsilon$ is the map of change of variables. Note that by the assumption of smoothness of L , we have that Ψ is smooth and $\det \Psi = 1$ on L . Now the first term of the right-hand side of (4.11) vanishes as $\epsilon \rightarrow 0$ since ω is bounded and $\mathcal{H}^2(\partial D_\epsilon) \rightarrow 0$. Integrating by parts the second term and using theorem 3.10, we obtain

$$\begin{aligned}
 &\sum_{k=4}^6 \int_0^{2\pi} \int_0^l \epsilon \int_0^l \frac{\partial \tilde{\omega}_{2k}}{\partial \tau}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) u_{k-3}(\epsilon, \theta, \tau) d\tau d\theta = \\
 &= \sum_{k=4}^6 \int_0^{2\pi} \int_0^l \epsilon \int_0^l \frac{\partial \tilde{\omega}_{2k}}{\partial \tau}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) \left(\frac{\theta b_{k-3}}{2\pi} + O(1) \right) d\tau d\theta \\
 &+ \sum_{k=4}^6 \int_0^{2\pi} \int_0^l \epsilon \int_0^l \sum_{h=1}^3 \frac{\partial \tilde{\omega}_{2k}}{\partial x_{3+h}}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) \frac{\partial u_h}{\partial \tau}(\epsilon, \theta, \tau) \left(\frac{\theta b_{k-3}}{2\pi} + O(1) \right) d\tau d\theta, \tag{4.12}
 \end{aligned}$$

where $|O(1)| \leq c$ as $\epsilon \rightarrow 0$ so that its absolute value can be estimated by $\epsilon c \|\partial \tilde{\omega}_{2k} / \partial x_i\|_{L^\infty}$ thanks to theorem 3.10, whereby this term vanishes as well as $\epsilon \rightarrow 0$.

As for the third term of (4.11), we first set

$$R(\epsilon) := - \sum_{k=4}^6 \int_0^{2\pi} \int_0^l \epsilon \Delta \tilde{\omega}_{3k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) \frac{\partial u_{k-3}}{\partial \sigma}(\epsilon, \theta, \tau) d\tau d\theta,$$

with, recalling that $u_{k-3}(0^+, \theta, \tau) = \lim_{\epsilon \rightarrow 0^+} u_{k-3}(\epsilon \rho, \theta, \tau) = ((\theta)/(2\pi))b_k + c$,

$$\Delta \tilde{\omega}_{3k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) := \tilde{\omega}_{3k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) - \tilde{\omega}_{3k}(\hat{\tau}, u_{k-3}(0^+, \theta, \tau)),$$

where $\hat{\tau} := (0, 0, \tau)$. Since $\partial / \partial \sigma = 1/\epsilon(\partial / \partial \theta)$, we obtain

$$\begin{aligned}
 &- \sum_{k=4}^6 \int_0^l \int_0^{2\pi} \tilde{\omega}_{3k}(\hat{\tau}, u_{k-3}(0^+, \theta, \tau)) \frac{\partial u_{k-3}}{\partial \theta}(\epsilon, \theta, \tau) d\theta d\tau + R(\epsilon) = \\
 &= - \sum_{k=4}^6 \int_0^l \tilde{\omega}_{3k} \left(\hat{\tau}, \frac{b_1 \theta}{2\pi} + c, \frac{b_2 \theta}{2\pi} + c, \frac{b_3 \theta}{2\pi} + c \right) u_{k-3}(\epsilon, \theta, \tau) \Big|_0^{\theta=2\pi} d\tau \\
 &+ \sum_{k=4}^6 \int_0^l \int_0^{2\pi} \frac{d}{d\theta} \tilde{\omega}_{3k} \left(\hat{\tau}, \frac{b_1 \theta}{2\pi} + c, \frac{b_2 \theta}{2\pi} + c, \frac{b_3 \theta}{2\pi} + c \right) u_{k-3}(\epsilon, \theta, \tau) d\theta d\tau + R(\epsilon). \tag{4.13}
 \end{aligned}$$

Using theorem 3.4, for some constant $c > 0$, we have

$$\begin{aligned}
 |R(\epsilon)| &\leq \sum_{k=4}^6 \int_0^{2\pi} \int_0^l \epsilon \|\Delta \tilde{\omega}_{3k}(x, u(x))\|_{L^\infty(\partial D_\epsilon)} \left| \frac{\partial u_{k-3}}{\partial \sigma}(\epsilon, \theta, \tau) \right| d\tau d\theta \\
 &\leq c \int_0^{2\pi} \|\Delta \tilde{\omega}_{3k}(x, u(x))\|_{L^\infty(\partial D_\epsilon)} d\theta \rightarrow 0,
 \end{aligned}$$

as $\epsilon \rightarrow 0$, by (3.6) and being $\tilde{\omega}_{3k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau))$ uniformly continuous at $\epsilon = 0$, again thanks to theorem 3.10 and the fact that L is compact. So that letting $\epsilon \rightarrow 0$ in (4.13), using theorem 3.10, integrating by parts again, and taking into account the periodicity of ω , (4.13) becomes

$$\begin{aligned}
 &-\sum_{k=4}^6 \int_0^l \int_0^{2\pi} \omega_{3k} \left(\hat{\tau}, \frac{b_1\theta}{2\pi} + c, \frac{b_2\theta}{2\pi} + c, \frac{b_3\theta}{2\pi} + c \right) \frac{b_{k-3}}{2\pi} d\theta d\tau \\
 &= -\frac{1}{2\pi} \int_0^l \int_0^{2\pi} \left\langle \omega \left(\hat{\tau}, \frac{b_1\theta}{2\pi}, \frac{b_2\theta}{2\pi}, \frac{b_3\theta}{2\pi} \right), \bar{\tau} \wedge \bar{b} \right\rangle d\theta d\tau \\
 &= \mathcal{L} \wedge \bar{b}(\omega),
 \end{aligned} \tag{4.14}$$

since in the local basis $\bar{\tau} = (\bar{\tau}, 0) = (0, 0, 1, 0, 0, 0)$. The proof is completed. □

4.3. Extensions of theorem 4.1

In this section, we discuss some extensions of theorem 4.1.

4.3.1. The case with a finite number of loops

THEOREM 4.5. *Let S be the union of $N > 0$ Lipschitz surfaces S_k in Ω whose boundary L is the union of the corresponding boundaries L_k , that is, closed curves in Ω . Let $b = (b_1, b_2, b_3) \in 2\pi\mathbb{Z}^3$ and let $u = (u_1, u_2, u_3) : \Omega \rightarrow \mathbb{R}^3$ be a map such that u_i satisfies (3.2) with $b = b_i, i = 1, 2, 3$. Then \mathcal{G}_u is an integral current in $\mathcal{D}_3(\Omega \times \mathbb{T}^3)$ and (4.4) holds.*

Proof. Let us first suppose that S_k and L_k are smooth and that the curves L_k are mutually disjoint. Then we will obtain the general result by approximation by means of theorem 2.3, arguing as in the proof of Corollary 4.3. Since N is finite, we see that L is compact and there is a tubular neighbourhood around L . We can then argue as in the proof of theorem 4.1, obtaining a formula similar to (4.13). The thesis follows since $R(\epsilon)$ in (4.13) still vanishes thanks to corollary 3.8, and $\partial u_{k-3}/\partial \sigma(\epsilon, \theta, \tau)$ in (4.12) still tends to 0. □

4.3.2. Perturbation by a smooth function

THEOREM 4.6. Let S, L, b and u as in theorem 4.5, and let $v \in C^1(\bar{\Omega}, \mathbb{R}^3)$. Then \mathcal{G}_{u+v} is the integral current in $\mathcal{D}_3(\Omega \times \mathbb{T}^3)$ given by

$$\partial \mathcal{G}_{u+v}(\omega) = \mathcal{L} \wedge \vec{b}(\omega) + \mathcal{C}_{u+v}(\omega), \tag{4.15}$$

for all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{T}^3)$, with \mathcal{C}_{u+v} defined as

$$\mathcal{C}_{u+v}(\omega) = \frac{1}{2\pi} \int_L \int_0^{2\pi} \left\langle \omega \left(x, \frac{b\theta}{2\pi} + v(x) \right), \frac{\partial \bar{v}}{\partial \tau} \wedge \bar{b} \right\rangle d\theta d\mathcal{H}^1(x), \tag{4.16}$$

for all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{T}^3)$ and with $\bar{v} = (0, v) \in C^1(\bar{\Omega}, \mathbb{R}^3 \times \mathbb{R}^3)$. In particular, it holds

$$M(\partial \mathcal{G}_{u+v}) \leq C(1 + \|Dv\|_{L^\infty(\Omega)}) |\mathcal{L} \otimes b|(\Omega). \tag{4.17}$$

Proof. As in theorem 4.1, we first prove the result for a smooth loop L and then we obtain the general case arguing as in theorem 4.5. Let us check that $u + v \in \mathcal{A}_p(\Omega, \mathbb{T}^3)$. To this aim let us prove that $\text{adj}(Du + Dv)$ and $\det(Du + Dv)$ are summable functions. Since the rows of Du are linearly dependent it follows that

$$\det(Du + Dv) = \det \begin{pmatrix} Dv_1 \\ Dv_2 \\ Dv_3 \end{pmatrix} + \det \begin{pmatrix} Du_1 \\ Dv_2 \\ Dv_3 \end{pmatrix} + \det \begin{pmatrix} Dv_1 \\ Du_2 \\ Dv_3 \end{pmatrix} + \det \begin{pmatrix} Dv_1 \\ Dv_2 \\ Du_3 \end{pmatrix}.$$

Since $Dv_i \in C^0(\bar{\Omega}, \mathbb{R}^3)$, in particular, it is bounded so that all the determinants belong to $L^p(\Omega, \mathbb{R}^3)$ thanks to (3.6). A similar argument applies for $\text{adj}(Du + Dv)$.

To compute the boundary of \mathcal{G}_{u+v} , we proceed as in the proof of Corollary 4.3 (and theorem 4.1), resulting in (4.10). This formula, setting $w := u + v$, takes the form

$$\begin{aligned} & \int_{\partial D_\epsilon} \omega_{23}(x, w(x)) + \sum_{k=1}^3 \left(\omega_{2k}(x, w(x)) \frac{\partial u_k}{\partial \tau}(x) - \omega_{3k}(x, w(x)) \frac{\partial u_k}{\partial \sigma}(x) \right) d\mathcal{H}^2(x) \\ & + \int_{\partial D_\epsilon} \sum_{k=1}^3 \left(\omega_{2k}(x, w(x)) \frac{\partial v_k}{\partial \tau}(x) - \omega_{3k}(x, w(x)) \frac{\partial v_k}{\partial \sigma}(x) \right) d\mathcal{H}^2(x) + \\ & + \sum_{4 \leq i < j \leq 6} \int_{\partial D_\epsilon} \omega_{ij}(x, w(x)) \tilde{M}_i^j(D(\text{Id} \times v) \lrcorner_{\partial D_\epsilon}(x)) d\mathcal{H}^2(x) \\ & + \sum_{4 \leq i \neq j \leq 6} \int_{\partial D_\epsilon} \omega_{ij}(x, w(x)) \left(\frac{\partial u_{i-3}}{\partial \sigma} \frac{\partial v_{j-3}}{\partial \tau} - \frac{\partial u_{i-3}}{\partial \tau} \frac{\partial v_{j-3}}{\partial \sigma} \right) d\mathcal{H}^2(x), \tag{4.18} \end{aligned}$$

where in the last term we have used the convention, with a little abuse of notation, that $\omega_{ji} := -\omega_{ij}$ for $i < j$. The first row, as seen by (4.11), tends to (4.14), the second and the third ones vanish as $\epsilon \rightarrow 0$ since v is smooth and its partial derivatives are bounded. The terms of the last row containing $\partial u_{i-3}/\partial \tau$ and $\partial v_{j-3}/\partial \sigma$ vanish

again thanks to theorem 3.10 and the smoothness of L . It remains to study the term

$$\begin{aligned} & \sum_{4 \leq i \neq j \leq 6} \int_{\partial D_\epsilon} \omega_{ij}(x, w(x)) \frac{\partial u_{i-3}}{\partial \sigma} \frac{\partial v_{j-3}}{\partial \tau} d\mathcal{H}^2(x) = \\ & = R^w(\epsilon) + \sum_{4 \leq i \neq j \leq 6} \int_0^l \int_0^{2\pi} \epsilon \tilde{\omega}_{ij}(\hat{\tau}, w(0^+, \theta, \tau)) \frac{\partial v_{j-3}}{\partial \tau}(\hat{\tau}) \frac{\partial u_{i-3}}{\partial \sigma}(\epsilon, \theta, \tau) d\theta d\tau, \end{aligned} \tag{4.19}$$

with $\hat{\tau} := (0, 0, \tau)$ and

$$R^w(\epsilon) = \sum_{4 \leq i \neq j \leq 6} \int_0^l \int_0^{2\pi} \epsilon \Delta \left(\omega_{ij} \frac{\partial v_{j-3}}{\partial \tau} \right) (\epsilon, \theta, \tau) \frac{\partial u_{i-3}}{\partial \sigma}(\epsilon, \theta, \tau) d\theta d\tau,$$

and

$$\begin{aligned} \Delta \left(\omega_{ij} \frac{\partial v_{j-3}}{\partial \tau} \right) (\epsilon, \theta, \tau) & := \tilde{\omega}_{ij}(\epsilon, \theta, \tau, w(\epsilon, \theta, \tau)) \frac{\partial v_{j-3}}{\partial \tau}(\epsilon, \theta, \tau) \\ & \quad - \tilde{\omega}_{ij}(\hat{\tau}, w(0^+, \theta, \tau)) \frac{\partial v_{j-3}}{\partial \tau}(\hat{\tau}). \end{aligned}$$

Arguing as for $R(\epsilon)$ in (4.13) we see that $R^w(\epsilon)$ is negligible as $\epsilon \rightarrow 0$, while arguing as in (4.13) and taking into account that $w(0^+, \theta, \tau) = (b\theta/2\pi) + c + v(\hat{\tau})$, the last term in (4.19) tends to

$$\begin{aligned} & \sum_{4 \leq i \neq j \leq 6} \int_0^l \int_0^{2\pi} \omega_{ij} \left(\hat{\tau}, \frac{b\theta}{2\pi} + c + v(\hat{\tau}) \right) \frac{\partial v_{j-3}}{\partial \tau}(\hat{\tau}) \frac{b_{i-3}}{2\pi} d\theta d\tau \\ & = \sum_{4 \leq i \neq j \leq 6} \int_0^l \int_0^{2\pi} \omega_{ij} \left(\hat{\tau}, \frac{b\theta}{2\pi} + v(\hat{\tau}) \right) \frac{\partial v_{j-3}}{\partial \tau}(\hat{\tau}) \frac{b_{i-3}}{2\pi} d\theta d\tau \\ & = \sum_{4 \leq i < j \leq 6} \int_0^l \int_0^{2\pi} \omega_{ij} \left(\hat{\tau}, \frac{b\theta}{2\pi} + v(\hat{\tau}) \right) \left(\frac{\partial v_{j-3}}{\partial \tau}(\hat{\tau}) \frac{b_{i-3}}{2\pi} - \frac{\partial v_{i-3}}{\partial \tau}(\hat{\tau}) \frac{b_{j-3}}{2\pi} \right) d\theta d\tau \\ & = \mathcal{C}_{u+v}(\omega), \end{aligned}$$

where we have considered the periodicity of ω . The bound (4.17) now readily follows. □

5. Main result second form: Euclidean setting and jump surface

In this section, we express the boundary of the graph of u as a current in $\Omega \times \mathbb{R}^3$, hence without avoiding the jump of u on the surface S .

5.1. Expression of the boundary graph for \mathbb{R}^3 -valued displacement

In this paper, the main result theorem 4.1 is presented without referring to the surface S (in the final formulae) since torus-valued maps are considered. Nonetheless, the formulae in the Euclidean setting might be of interest with a view to the

introduction of misfit energy terms as explained in § 6.6. The derivation of such formulae is rather technical, though they do not require new notions with respect to those exposed in the paper core. This is the reason why the complete proof is given in Appendix A.

Let us introduce the following notation. For all $b \in \mathbb{R}^3$ and all $r \in \mathbb{R}^3$, we define the 1-current $\vec{b}_r \in \mathcal{D}_1(\mathbb{R}^3)$ as

$$\vec{b}_r(\omega) := -\frac{1}{2\pi} \int_0^{2\pi} \left\langle \omega \left(r_1 + \frac{b_1\theta}{2\pi}, r_2 + \frac{b_2\theta}{2\pi}, r_3 + \frac{b_3\theta}{2\pi} \right), b \right\rangle d\theta, \tag{5.1}$$

for any 1-form $\omega \in \mathcal{D}^1(\mathbb{R}^3)$. Moreover, $M(\vec{b}_r) = |b|$ for all $r \in \mathbb{R}^3$.

Let τ be an arc length parameter along the curve L . As in theorem 3.10, we will denote by $u(\tau)$ the value of the limit as $\epsilon \rightarrow 0$ of $u(\epsilon, \theta = 0, \tau)$, with $\{\epsilon, \theta, \tau\}$ a system of cylindrical coordinates around L taken in such a way that $\theta = 0$ corresponds to the set of points in the surface S , so that the lower and upper traces of u on S can be denoted by $u(\epsilon, 0, \tau)$ and $u(\epsilon, 2\pi, \tau)$, respectively (at least in a small neighbourhood of L). Since the jump of u is the constant b on S we can assume $u(\epsilon, 2\pi, \tau) = b + u(\epsilon, 0, \tau)$. With this convention $u(\tau) := \lim_{\epsilon \rightarrow 0} u(\epsilon, 0, \tau)$.

We have denoted by $\mathcal{L} \wedge \vec{b}_u$ the 2-current in $\Omega \times \mathbb{R}^3$ defined as

$$\mathcal{L} \wedge \vec{b}_u(\omega) = -\frac{1}{2\pi} \int_L \int_0^{2\pi} \left\langle \omega \left(x, u(\tau) + \frac{b\theta}{2\pi} \right), \bar{\tau} \wedge \bar{b} \right\rangle d\theta d\mathcal{H}^1(x), \tag{5.2}$$

for any 2-form $\omega \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$.

THEOREM 5.1. *Let L be a smooth closed curve in Ω and let S be a smooth surface with $\partial S = L$. Let $b \in \mathbb{R}^3$ and let $u_i \in SBV(\Omega)$ be the solution to (3.1) with $b = b_i$ for $i = 1, 2, 3$. Let $v \in \mathcal{C}^1(\bar{\Omega}, \mathbb{R}^3)$ and set $u = (u_1, u_2, u_3)$, and $w := u + v$. Then \mathcal{G}_u and \mathcal{G}_w are integral currents in $\mathcal{D}_3(\Omega \times \mathbb{R}^3)$ and it holds*

$$\begin{aligned} \partial \mathcal{G}_u(\omega) &= \mathcal{L} \wedge \vec{b}_u(\omega) + \int_S (\omega_{12}(x, u(x) + b) - \omega_{12}(x, u(x))) d\mathcal{H}^2(x) \\ &\quad + \sum_{k=4}^6 \int_S b_{k-3} (\omega_{1k}(x, u(x) + b) - \omega_{1k}(x, u(x))) \frac{\partial \hat{u}}{\partial \tau_2} d\mathcal{H}^2(x) \\ &\quad - \sum_{k=4}^6 \int_S b_{k-3} (\omega_{2k}(x, u(x) + b) - \omega_{2k}(x, u(x))) \frac{\partial \hat{u}}{\partial \tau_1} d\mathcal{H}^2(x), \end{aligned} \tag{5.3}$$

for all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$, where $\{\tau_1, \tau_2\}$ is an orthogonal basis for S , and \hat{u} is the solution to (3.1) with $b = 1$. In the previous formula u and $u + b$ are the two traces of u on the lower and upper face of S . Moreover,

$$\begin{aligned} \partial \mathcal{G}_w(\omega) &= \sum_{4 \leq i < j \leq 6} \int_S (\omega_{ij}(x, w(x) + b) - \omega_{ij}(x, w(x))) \\ &\quad \left(\frac{\partial w_{i-3}}{\partial \tau_1} \frac{\partial w_{j-3}}{\partial \tau_2} - \frac{\partial w_{j-3}}{\partial \tau_1} \frac{\partial w_{i-3}}{\partial \tau_2} \right) d\mathcal{H}^2(x) \\ &\quad + \int_S (\omega_{12}(x, w(x) + b) - \omega_{12}(x, w(x))) d\mathcal{H}^2(x) \end{aligned}$$

$$\begin{aligned}
 & + \sum_{k=4}^6 \int_S (\omega_{1k}(x, w(x) + b) - \omega_{1k}(x, w(x))) \frac{\partial w_{k-3}}{\partial \tau_2} d\mathcal{H}^2(x) \\
 & - \sum_{k=4}^6 \int_S (\omega_{2k}(x, w(x) + b) - \omega_{2k}(x, w(x))) \frac{\partial w_{k-3}}{\partial \tau_1} d\mathcal{H}^2(x) \\
 & + \mathcal{L} \wedge \vec{b}_w(\omega) + \mathcal{C}_w(\omega),
 \end{aligned} \tag{5.4}$$

for all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$, where \mathcal{C}_w is a rectifiable 2-current with support in $C \times \mathbb{R}^3$ given by

$$\mathcal{C}_w(\omega) := \frac{1}{2\pi} \int_L \int_0^{2\pi} \left\langle \omega \left(x, \frac{b\theta}{2\pi} + w(x) \right), \frac{\partial \bar{v}}{\partial \tau}(x) \wedge \bar{b} \right\rangle d\theta d\mathcal{H}^1(x), \tag{5.5}$$

with $\bar{v} = (0, v)$ and for all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$. Furthermore,

$$M(\mathcal{C}_w) \leq |\mathcal{L} \otimes b|(\Omega) \|Dv\|_{L^\infty}.$$

In particular, there exists a constant $c > 0$ depending only on the geometry of L such that

$$M(\partial\mathcal{G}_w) \leq |\mathcal{L} \otimes b|(\Omega) + c\mathcal{H}^2(S), \tag{5.6}$$

and

$$M(\partial\mathcal{G}_w) \leq (1 + \|Dv\|_{L^\infty})(|\mathcal{L} \otimes b|(\Omega) + c\mathcal{H}^2(S)) + \mathcal{H}^2(S) \|Dv\|_{L^\infty}^2. \tag{5.7}$$

For a proof, we refer to Appendix A.

6. Application to dislocations at the continuum scale

6.1. Preliminaries on dislocations at the continuum scale

In this section, we will often identify loops with 1 integral currents so that we will employ the same notation \mathcal{L} for both. The following lemma characterizes the divergence-free measures which are supported on a 1-dimensional set as dislocation measures.

LEMMA 6.1 (Dislocation measure). *Let $\mu \in \mathcal{M}_b(\Omega, \mathbb{R}^{3 \times 3})$ be a divergence-free measure which is absolutely continuous with respect to the \mathcal{H}^1 -measure restricted to a simple Lipschitz curve \mathcal{L} with tangent vector τ and such that \mathcal{L} is either closed or ends at the boundary of Ω . Then μ is a dislocation-measure, that is, there exists a constant vector b such that*

$$\mu = b \otimes \tau \mathcal{H}^1_{\llcorner \mathcal{L}}.$$

Proof. By definition $\langle \mu, \varphi \rangle = \int_{\mathcal{L}} M(x) \cdot \varphi(x) d\mathcal{H}^1(x)$ for every $\varphi \in \mathcal{C}_c^\infty(\Omega, \mathbb{R}^{3 \times 3})$, with $M \in L^1(\mathcal{L}, \mathbb{R}^{3 \times 3})$. Moreover, $\langle \mu, D\psi \rangle = 0$ for every $\psi \in \mathcal{C}_c^\infty(\Omega, \mathbb{R}^3)$. Let $\{\nu, \sigma, \tau\}$ be a local orthogonal basis attached to $x \in \mathcal{L}$, and let (t, z, s) be the associated coordinate system in a neighbourhood of \mathcal{L} . By orthogonal decomposition, $M_{ij} = M_{ik}\tau_k\tau_j + M_{ik}\nu_k\nu_j + M_{ik}\sigma_k\sigma_j$ and $\varphi_{ij} = \varphi_{ik}\tau_k\tau_j + \varphi_{ik}\nu_k\nu_j + \varphi_{ik}\sigma_k\sigma_j$

for all $i = 1, 2, 3$. Let $\xi, \eta : \mathcal{L} \rightarrow \mathbb{R}^3$ be smooth functions with compact support and set $\psi(t, z, s) = \xi(s)z + \eta(s)t$. One has $\partial_s \psi(x) = \dot{\xi}z + \dot{\eta}t = 0$ for $x = (0, 0, s) \in \mathcal{L}$. Moreover, $\partial_t \psi|_{\mathcal{L}} = \eta$ and $\partial_z \psi|_{\mathcal{L}} = \xi$, so that one has $0 = \int_{\mathcal{L}} M(x) \cdot D\psi(x) d\mathcal{H}^1 = \int_{\mathcal{L}} (M_{ij}\eta_i\nu_j + M_{ij}\xi_i\sigma_j) d\mathcal{H}^1$, and hence $M_{ij} = b_i\tau_j$ with $b_i := M_{ik}\tau_k$. Taking now $\varphi = D\psi$ for an arbitrary ψ smooth with compact support, it results from the closedness property of \mathcal{L} that $0 = \int_{\mathcal{L}} b \cdot \partial_\tau \psi d\mathcal{H}^1 = - \int_{\mathcal{L}} \partial_\tau b \cdot \psi d\mathcal{H}^1$, and b is constant. \square

According to lemma 6.1, a dislocation loop is defined as a simple closed curve \mathcal{L} in Ω which has an associated Burgers vector $b \in \mathbb{Z}^3$. The deformation gradient F around \mathcal{L} satisfies the condition

$$- \text{Curl } F = \Lambda_{\mathcal{L}^b}^T := b \otimes \tau \mathcal{H}^1 \llcorner_{\mathcal{L}},$$

where τ is an oriented tangent vector to \mathcal{L} . For any $b \in 2\pi\mathbb{Z}^3$ we call a b -dislocation current a 1-integer multiplicity current \mathcal{L}^b that produces a curl of the deformation gradient given by the density $\Lambda_{\mathcal{L}^b}$, hence satisfying

$$\langle \Lambda_{\mathcal{L}^b}, w \rangle = \mathcal{L}^b((wb)^*), \tag{6.1}$$

for every $w \in C_c^\infty(\Omega, \mathbb{R}^{3 \times 3})$, where in the right-hand side $(wb)^*$ is the covector writing $(wb)^* := w_{kj}b_j dx^k$ (with sums on the repeated indices). Moreover, $\Lambda_{\mathcal{L}^b}$ is a Radon measure as soon as $M(\mathcal{L}^b)$ is finite. In the sequel, we will use the following shortcut notation:

$$\Lambda_{\mathcal{L}^b} = \mathcal{L}^b \otimes b = \tau^b \otimes b \theta^b \mathcal{H}^1 \llcorner_{\mathcal{L}}, \tag{6.2}$$

with θ^b the multiplicity of the vector b (see [37] for details).

DEFINITION 6.2 (Regular dislocation). Let us set $\mathcal{B} := 2\pi\mathbb{Z}^3$ the set of admissible Burgers vectors. A *regular dislocation* is a sequence of b -dislocation currents $\mathcal{L} := \{\mathcal{L}^b\}_{b \in \mathcal{B}}$. We associate to each dislocation a *dislocation current*, still denoted by \mathcal{L} , and the associated *dislocation density* $\Lambda_{\mathcal{L}}$,

$$\mathcal{L} := \sum_{b \in \mathcal{B}} \mathcal{L}^b, \quad \Lambda_{\mathcal{L}} := \sum_{b \in \mathcal{B}} \Lambda_{\mathcal{L}^b}. \tag{6.3}$$

With this definition it is possible to model dislocations with every possible Burgers vector. It is possible to split the current \mathcal{L} on the canonical basis of \mathbb{R}^3 ,

$$\mathcal{L} = \mathcal{L}_1 + \mathcal{L}_2 + \mathcal{L}_3,$$

in such a way that \mathcal{L}_i has e_i as associated Burgers vector and satisfies

$$\Lambda_{\mathcal{L}} = \Lambda_{\mathcal{L}_1} + \Lambda_{\mathcal{L}_2} + \Lambda_{\mathcal{L}_3} = \sum_{i=1}^3 \mathcal{L}_i \otimes e_i.$$

Moreover, as proved in [37], one has

$$|\mathcal{L}_i|_\Omega \leq c |\Lambda_{\mathcal{L}}|_{\mathcal{M}(\Omega)}, \tag{6.4}$$

for some constant $c > 0$ independent of i and Ω .

6.2. Invertibility of the curl

Let $\mathcal{M}_{\text{div}}(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ denote the space of divergence-free bounded Radon measures with values in $\mathbb{R}^{3 \times 3}$. The following result is an extension of a result by Bourgain and Brezis [9] that we can prove thanks to the results of previous sections.

THEOREM 6.3 (Biot-Savart). *Let $\hat{\Omega} \subset \mathbb{R}^3$ be an open simply connected and smooth set. Let μ be a tensor-valued Radon measure such that $\mu \in \mathcal{M}_{\text{div}}(\hat{\Omega}, \mathbb{R}^{3 \times 3})$. Then there exists a unique $F \in L^1(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ solution of*

$$\begin{cases} -\text{Curl } F &= \mu & \text{in } \hat{\Omega} \\ \text{div } F &= 0 & \text{in } \hat{\Omega} \\ FN &= 0 & \text{on } \partial\hat{\Omega}. \end{cases} \tag{6.5}$$

Moreover, F belongs to $L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ for all p with $1 \leq p < 3/2$ and for all such p there exists a constant $C > 0$ satisfying

$$\|F\|_p \leq C|\mu|(\hat{\Omega}). \tag{6.6}$$

In the case that $\mu = b \otimes \tau \mathcal{H}^1 \llcorner \mathcal{L}$, for some $b \in \mathbb{R}^3$ and a \mathcal{C}^2 -closed curve \mathcal{L} in $\hat{\Omega}$ with unit oriented tangent vector τ , then the solution F belongs to $L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ for all $p < 2$.

The proof of the first statement is a straightforward extensions of the main results in [9], also discussed in [15]. Thus we omit it. We address here only the proof of the second statement.

Proof of the sharp result for a smooth curve. We will use the fact that the solution F is a smooth perturbation of the field G satisfying

$\begin{cases} -\text{Curl } G &= b \otimes \tau \mathcal{H}^1 \llcorner \mathcal{L} & \text{in } \hat{\Omega} \\ \text{div } G &= 0 & \text{in } \hat{\Omega}. \end{cases}$, and given explicitly by ∇u with u given by (3.2) (see theorem 3.13). Now, we will use the estimate (3.6) for G , proven in lemma 3.4, which asserts that there is a constant $c > 0$ depending only on the curvature of \mathcal{L} such that, for $x \in \Omega$,

$$|\nabla G|(x) \leq c|b|l \left(1 + \frac{1}{d(x, \mathcal{L})} \right), \tag{6.7}$$

where l denotes the length of the curve \mathcal{L} . Since L has bounded curvature there exists $\delta_0 > 0$ and a constant $\tilde{c} > 0$ such that there holds

$$\forall \delta < \delta_0 \quad \mathcal{H}^2(\{x : d(x, \mathcal{L}) = \delta\}) \leq 2\pi\tilde{c}\delta. \tag{6.8}$$

Using (6.7), (6.8), exploiting that the map $x \mapsto d(x, \mathcal{L})$ has gradient such that $|\nabla d| = 1$ a.e. in Ω , and employing the co-area formula, we estimate

$$\int_{\hat{\Omega}} |G|^p dx = \int_0^{\text{diam}(\hat{\Omega})} \int_{\{d=r\}} |G|^p d\mathcal{H}^2 dr \leq c + c \int_0^{\text{diam}(\hat{\Omega})} \frac{r}{r^p} dr < +\infty, \tag{6.9}$$

whenever $p < 2$. □

6.3. Functional properties

Let $1 \leq p < \infty$ and introduce the vector space of tensor-valued fields

$$\mathcal{BC}^p(\Omega, \mathbb{R}^{3 \times 3}) := \{F \in L^p(\Omega, \mathbb{R}^{3 \times 3}) \text{ s.t. } \text{Curl } F \in \mathcal{M}_b(\bar{\Omega}, \mathbb{R}^{3 \times 3})\}, \tag{6.10}$$

which endowed with the norm $\|F\|_{\mathcal{BC}^p} := \|F\|_p + |\text{Curl } F|(\bar{\Omega})$, is a Banach space. Let us define

$$L^p_{\text{div}}(\Omega, \mathbb{R}^{3 \times 3}) := \{F \in L^p(\Omega) \text{ s.t. } \text{div } F = 0\},$$

and the space

$$\tilde{\mathcal{V}}^p(\Omega) := \{V \in L^p_{\text{div}}(\Omega, \mathbb{R}^{3 \times 3}) \text{ s.t. } \text{Curl } V \in L^p(\Omega, \mathbb{R}^{3 \times 3}), VN = 0 \text{ on } \partial\Omega\}. \tag{6.11}$$

With a view to applications, it is convenient to define another open set $\hat{\Omega}$ such that $\Omega \subset\subset \hat{\Omega}$. We will also assume that both Ω and $\hat{\Omega}$ are simply connected. In addition, we assume that $\hat{\Omega}$ has a smooth boundary.

Let $1 < p < 2$ and let $F \in \mathcal{BC}^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ be such that, by virtue of theorem 6.3, $-\text{Curl } F = b \otimes \mathcal{L}$, with $b \in 2\pi\mathbb{Z}^3$ and \mathcal{L} a 1-integer multiplicity current which is closed and with compact support in $\hat{\Omega}$. The Helmholtz decomposition in $L^p(\hat{\Omega}, \mathbb{R}^3)$ provides $v \in W^{1,p}(\hat{\Omega}, \mathbb{R}^3)$ and $G \in \tilde{\mathcal{V}}^p(\hat{\Omega})$ such that (see [28] for details and further references)

$$F = Dv + \text{Curl } G. \tag{6.12}$$

If we set $V := \text{Curl } G$, then $\text{div } V = 0$, while since $-\text{Curl } F = b \otimes \mathcal{L}$, we also have $-\text{Curl } V = b \otimes \mathcal{L}$. Thanks to the decomposition theorem for 1-integer multiplicity currents (theorem 2.1) we find a sequence of Lipschitz maps

$$f_k : S^1 \rightarrow \hat{\Omega} \text{ such that } \mathcal{L} = \sum_{k>0} f_{k\#} \llbracket S^1 \rrbracket. \tag{6.13}$$

Let us denote by \mathcal{L}_k the closed Lipschitz curves $f_k(S^1)$.

6.4. Expression of theorem 4.1 for a finite number of dislocation loops

THEOREM 6.4. *Let $b \in 2\pi\mathbb{Z}^3$ be fixed, $1 < p < 2$, \mathcal{L} be a closed integral current with compact support in $\hat{\Omega}$, and let $V \in \mathcal{BC}^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}) \cap L^p_{\text{div}}(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ be such that $-\text{Curl } V = b \otimes \mathcal{L}$ in $\hat{\Omega}$. Then there exists a map $\tilde{u} \in \mathcal{A}_p(\hat{\Omega}, \mathbb{T}^3)$ such that $\nabla \tilde{u} = V$ almost everywhere in $\hat{\Omega}$, and*

$$M(\partial\mathcal{G}_{\tilde{u}}) \leq c|\mathcal{L} \otimes b|(\hat{\Omega})(1 + |\mathcal{L} \otimes b|(\hat{\Omega})), \tag{6.14}$$

with $c > 0$ a constant depending only on $\hat{\Omega}$. Moreover, $\tilde{u} = u - v$ with $v \in \mathcal{C}^1(\bar{\hat{\Omega}}, \mathbb{R}^3)$, $u \in \mathcal{A}_p(\hat{\Omega}, \mathbb{T}^3)$, and

$$\partial\mathcal{G}_u(\omega) = \mathcal{L} \wedge \vec{b}(\omega) = \Lambda_{\mathcal{L}}(\varphi^\omega) := \mathcal{L} \otimes b(\varphi^\omega), \tag{6.15}$$

for all $\omega \in \mathcal{D}^2(\hat{\Omega} \times \mathbb{T}^3)$.

Recalling remark 4.2, φ^ω is given by (4.5). Hence, theorem 6.4 provides an expression of the dislocation density in terms of the boundary of the graph of the torus-valued displacement field u .

Proof. Let us first assume $p < 3/2$. To prove the Theorem, we will use the fact that the following system

$$\begin{cases} -\operatorname{Curl} U &= \mu & \text{in } \hat{\Omega} \\ \operatorname{div} U &= 0 & \text{in } \hat{\Omega} \\ UN &= 0 & \text{on } \partial\hat{\Omega}, \end{cases} \tag{6.16}$$

has by theorem 6.3 a unique solution that also satisfies $\|U\|_{L^p} \leq c|\mu|(\hat{\Omega})$, with $c = c(\hat{\Omega})$. Another key fact is the following: if $\hat{\Omega}$ is a bounded open set with smooth boundary, $g \in \mathcal{C}^{1,\alpha}(\partial\hat{\Omega}, \mathbb{R}^3)$ with $\int_{\partial\hat{\Omega}} g d\mathcal{H}^2 = 0$, and $v \in \mathcal{C}^{2,\alpha}(\hat{\Omega}, \mathbb{R}^3)$ is the zero-mean-value solution to

$$\begin{cases} \Delta v &= 0 & \text{in } \hat{\Omega} \\ \partial_N v &= g & \text{on } \partial\hat{\Omega}, \end{cases} \tag{6.17}$$

then $\|v\|_{\mathcal{C}^{2,\alpha}} \leq c\|g\|_{\mathcal{C}^{1,\alpha}(\partial\hat{\Omega})}$, with $c = c(\hat{\Omega})$ (see, e.g., [22, theorems 6.30 and 6.31]).

We use the decomposition (6.13) for \mathcal{L} and we first suppose that the maps f_k are smooth. The general case will follow using an approximation argument and proceeding as in the proof of theorem 4.5. If L_k is a smooth closed curve, we can choose a smooth surface S_k with boundary L_k , according to assumption 3.1. Then we set $S := \cup_k S_k$ and $L := \cup_k L_k$, we seek a solution u of (3.1) with these S and L . Let us also set $\hat{S}_n := \cup_{k=0}^n S_k$ and $\hat{L}_n := \cup_{k=0}^n L_k$. For $i = 1, 2, 3$, let u_i^n be the solution of (3.1) with \hat{S}_n , \hat{L}_n , and b_i . Theorem 3.13 and remark 3.15 show that the distributional divergence of ∇u^n is zero, while the curl is given by $-b \otimes \sum_{k=0}^n f_{k\#} \llbracket S^1 \rrbracket$. Up to subtracting a constant to u^n , we also suppose it has zero mean value.

By hypotheses, it holds $\inf_k d(L_k, \partial\hat{\Omega}) > 0$, and then u^n are of class \mathcal{C}^∞ on $\partial\hat{\Omega}$, and their $\mathcal{C}^h(\partial\hat{\Omega})$ norms are uniformly bounded with respect to n for all $h > 0$ (taking into account that the set $L = \cup_k L_k$ has finite length, and then $S = \cup_k S_k$ has finite \mathcal{H}^2 -measure). Let v^n be the solution to (6.17) with $g := \partial_N u^n$. From the estimates of this solution we find $\|v^n\|_{\mathcal{C}^1} \leq c_1 \|\partial_N u^n\|_{\mathcal{C}^{1,\alpha}(\partial\hat{\Omega})} < c_2$, for some constant c_2 independent of n . Setting $\tilde{u}^n := u^n - v^n$, we see that $\nabla \tilde{u}^n$ solves system (6.5) with $\mu = \mu^n := b \otimes \sum_{k=0}^n f_{k\#} \llbracket S^1 \rrbracket$, so that we also have $\|\nabla \tilde{u}^n\|_p \leq |\mu^n|(\hat{\Omega}) < c_3$, with c_3 independent of n . In particular, we get $\|u^n\|_{W^{1,p}} \leq \|v^n\|_{W^{1,p}} + \|\tilde{u}^n\|_{W^{1,p}} \leq c$, for a constant $c > 0$ independent of n . Therefore, up to subsequences, $u^n \rightharpoonup u$ weakly in $W^{1,p}(\hat{\Omega}, \mathbb{R}^3)$, for some $u \in W^{1,p}(\hat{\Omega}, \mathbb{R}^3)$. Similarly, $\tilde{u}^n \rightharpoonup \tilde{u}$ and $v^n \rightharpoonup v$ weakly in $W^{1,p}(\hat{\Omega}, \mathbb{R}^3)$, and from $u^n = \tilde{u}^n + v^n$ we infer $u = \tilde{u} + v$. Since the rows of ∇u^n are equal up to a multiplicative factor, we also get that all the minors of u^n are uniformly bounded in L^p . Then, by theorem 2.3 and lemma 4.4, u^n weakly converge in $\mathcal{A}_p(\hat{\Omega}, \mathbb{T}^3)$ to u (up to extracting further subsequences). Moreover, theorem 4.5 implies that for every $n > 0$ equation (4.4) holds for u^n , with \mathcal{L} replaced by $\sum_{k=0}^n f_{k\#} \llbracket S^1 \rrbracket$. Now, lemma 2.5 implies that \mathcal{G}_u is an integral current whose

boundary satisfies

$$\partial\mathcal{G}_u(\omega) = \sum_{k=1}^{\infty} f_{k\sharp} \llbracket S^1 \rrbracket \wedge \vec{b}(\omega), \tag{6.18}$$

for all $\omega \in \mathcal{D}^2(\hat{\Omega} \times \mathbb{T}^3)$. To conclude the proof it suffices to observe that the maps \tilde{u}^n are smooth in a neighbourhood of $\partial\hat{\Omega}$ with $\partial_N \tilde{u}^n$ vanishing on $\partial\hat{\Omega}$, and hence $\partial_N \tilde{u}$ also vanishes, in such a way that $\nabla \tilde{u}$ satisfies (6.5) with $\mu := b \otimes \mathcal{L}$. By the smoothness properties of v^n , it is also true that v satisfies (6.17) with a bounded and smooth $g = \partial_N u$, so it is smooth in $\hat{\Omega}$ and lemma 4.6 implies (4.17). We now compute g by using formula (3.3) and $d(L_k, \partial\hat{\Omega}) > 0$. Arguing similarly as in theorem 3.4 (actually deriving expression (3.3) and using the fact that $d(L, \partial\hat{\Omega}) > 0$) we show that there is a constant $c_4 > 0$ such that $\|g\|_{C^{1,\alpha}(\partial\hat{\Omega})} \leq \|g\|_{C^2(\partial\hat{\Omega})} \leq c_4 |b \otimes \mathcal{L}|(\hat{\Omega})$, so that the inequality $\|v\|_{C^1} \leq c \|g\|_{C^{1,\alpha}(\partial\hat{\Omega})}$ together with (4.17) gives (6.14). The thesis is proved when $p < 3/2$. Assume $V \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with $3/2 \leq p < 2$. In particular, $V \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with $p < 3/2$, so the previous argument shows that $V = \nabla \tilde{u} = \nabla u - \nabla v$ with \tilde{u} satisfying (6.14), u satisfying (6.15), and v of class C^1 . In particular, $\nabla u \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with $3/2 \leq p < 2$, and the thesis follows. \square

REMARK 6.5. By definition of the u^k , we have observed that for all k the three components u_i^k , $i = 1, 2, 3$, differ by a multiplicative factor. In particular, we have seen that their gradients ∇u_i^k (i.e., the rows of the matrix ∇u^k) are linearly dependent. As a consequence, the same is true for the gradients ∇u_i . Thus, the three components of the harmonic function v have as boundary data $\partial_N u_i$ three linearly dependent vector fields. This implies, by the uniqueness of solution of elliptic equations that also ∇v_i are linearly dependent and hence that the final matrix $V = \nabla u = \nabla \tilde{u} - \nabla v$ has linearly dependent rows. Therefore, its pointwise adjunct and determinant are constantly zero.

6.5. Two examples

In this section, we show how theorem 4.1 and its corollaries can be applied to obtain closure properties of suitable classes of deformation fields. This goal is achieved by means of two examples of variational problems. In both the examples we deal with two bounded, simply connected and smooth open sets Ω and $\hat{\Omega}$, with $\Omega \subset\subset \hat{\Omega}$. We fix a boundary datum α for the dislocation current in $\hat{\Omega} \setminus \Omega$ and a map $\hat{F} \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with $-\text{Curl } \hat{F} = \Lambda_\alpha^T$ on $\hat{\Omega}$ and satisfying some suitable properties. Then we are interested in the minimization problem

$$\min_{F \in \mathcal{F}} \mathcal{W}(F),$$

where F varies in a suitable class of admissible fields \mathcal{F} satisfying $F = \hat{F}$ on $\hat{\Omega} \setminus \Omega$.

6.5.1. *A variational problem related to dislocations in finite elasticity.* We deal with an energy \mathcal{W} with the form

$$\mathcal{W}(F) := \mathcal{W}_e(F) + \mathcal{W}_{\text{defect}}(\text{Curl } F), \tag{6.19}$$

where we assume the following properties on \mathcal{W}_e and $\mathcal{W}_{\text{defect}}$:

- (i) The following coerciveness condition holds: there exists positive constants $\alpha_0, \alpha_1, \beta_0, \beta_1$, and $p \geq 1$ such that

$$\mathcal{W}_e(F) \geq \beta_1(\|F\|_{L^p}^p + \|\text{adj} F\|_{L^p}^p + \|\det F\|_{L^p}^p) - \beta_0, \tag{6.20}$$

$$\mathcal{W}_{\text{defect}}(\Lambda) \geq \alpha_1|\Lambda|(\Omega) - \alpha_0. \tag{6.21}$$

- (ii) $\mathcal{W}_{\text{defect}}$ is a function on $\mathcal{M}_b(\Omega, \mathbb{R}^{3 \times 3})$ which is lower semicontinuous with respect to the weak* convergence.
- (iii) \mathcal{W}_e is a function of $M(F)$ (i.e., of $F, \text{adj} F$, and $\det F$), and it is lower semicontinuous in $M(F)$ with respect to the weak convergence in L^p .

We assume that $\hat{F} \in \mathcal{A}^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with $-\text{Curl } \hat{F} = \Lambda_\alpha^T$ on $\hat{\Omega}$, and the additional property that $\text{div } \hat{F} = 0$ on $\hat{\Omega}$. Let $b \in 2\pi\mathbb{Z}^3$ be a fixed Burgers vector, let $p > 1$, then the class of admissible deformation tensors is

$$\begin{aligned} \mathcal{F}_b^p := \{F \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}) : -\text{div } F = 0, -\text{Curl } F = b \otimes \mathcal{L} \\ \text{for some closed integral 1-current } \mathcal{L}, \text{ and } F = \hat{F} \text{ on } \hat{\Omega} \setminus \Omega\}. \end{aligned} \tag{6.22}$$

The minimization problem reads

$$F \in \underset{F \in \mathcal{F}_b^p}{\text{argmin}} \mathcal{W}. \tag{6.23}$$

The existence of a solution is provided by the following:

THEOREM 6.6. *Let $1 < p < 2$. If \mathcal{W} satisfies (i), (ii), and (iii), then there exists a solution $F \in \mathcal{F}_b^p$ of (6.23).*

Proof. We will apply the direct method. Let $\{F_k\}_{k>0}$ be a minimizing sequence. From the coerciveness (i), we see that there exist $F \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}), A \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}),$ and $D \in L^p(\hat{\Omega}),$ such that

$$F_k \rightharpoonup F \text{ weakly in } L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}), \tag{6.24a}$$

$$\text{adj} F_k \rightharpoonup A \text{ weakly in } L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}), \tag{6.24b}$$

$$\det F_k \rightharpoonup D \text{ weakly in } L^p(\hat{\Omega}), \tag{6.24c}$$

Moreover, we find a measure $\Lambda \in \mathcal{M}_b(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with

$$\Lambda_k \rightharpoonup \Lambda \text{ weakly* in } \mathcal{M}_b(\hat{\Omega}, \mathbb{R}^{3 \times 3}), \tag{6.25}$$

where we have set $\Lambda_k = \Lambda_{\mathcal{L}_k} = -(\text{Curl } F_k)^T$. As [37, lemma 7.5] shows, there exists a regular dislocation current \mathcal{L} such that $(\mathcal{L}_k)_i \rightharpoonup \mathcal{L}_i$ in $\mathcal{D}_1(\hat{\Omega})$ and $\Lambda = \Lambda_{\mathcal{L}} = \sum_{i=1}^3 \mathcal{L}_i \otimes e_i$. In order to prove the theorem, we have to show that $\text{div } F = 0, -\text{Curl } F = \Lambda_{\mathcal{L}}, A = \text{adj} F,$ and $D = \det F$.

The functions F_k satisfy the hypotheses of theorem 6.4, thus there exists a sequence of harmonic functions $\tilde{u}_k \in \mathcal{A}_p(\hat{\Omega}, \mathbb{T}^3)$ such that $\nabla \tilde{u}_k = F_k$ satisfying

$$M(\partial \mathcal{G}_{\tilde{u}_k}) \leq C|\mathcal{L}_k \otimes b|(\hat{\Omega})(1 + |\mathcal{L}_k \otimes b|(\hat{\Omega})) \leq M, \tag{6.26}$$

for some constant $M > 0$. By (6.24a) there exists $\tilde{u} \in \mathcal{A}_p(\hat{\Omega}, \mathbb{T}^3)$ such that, up to a subsequence,

$$\tilde{u}_k \rightarrow \tilde{u} \text{ weakly in } W^{1,p}(\hat{\Omega}, \mathbb{T}^3). \tag{6.27}$$

Now, thanks to (6.26), theorem 2.3 applies, obtaining

$$\text{adj}(\nabla \tilde{u}_k) \rightarrow \text{adj}(\nabla \tilde{u}) \text{ weakly in } L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}), \tag{6.28}$$

$$\det(\nabla \tilde{u}_k) \rightarrow \det(\nabla \tilde{u}) \text{ weakly in } L^p(\hat{\Omega}). \tag{6.29}$$

These read $\text{adj}F_k \rightharpoonup \text{adj}F$ weakly in $L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ and $\det F_k \rightharpoonup \det F$ weakly in $L^p(\hat{\Omega})$. Therefore, $A = \text{adj}F$ by (6.24b), and $\det F = D$ from (6.24c). Moreover, for every test function $\varphi \in C_c^\infty(\hat{\Omega}, \mathbb{R}^3)$, we have

$$\langle F, \nabla \varphi \rangle = \langle \nabla w, \nabla \varphi \rangle = \lim_{k \rightarrow \infty} \langle \nabla w_k, \nabla \varphi \rangle = \lim_{k \rightarrow \infty} \langle \text{div } F_k, \varphi \rangle = 0,$$

and it follows $\text{div } F = 0$ on $\hat{\Omega}$. Finally, we write

$$\langle \text{Curl } F_k, \varphi \rangle = \langle F_k, \text{Curl } \varphi \rangle \rightarrow \langle F, \text{Curl } \varphi \rangle = \langle \text{Curl } F, \varphi \rangle,$$

for all $\varphi \in C_c^\infty(\hat{\Omega}, \mathbb{R}^{3 \times 3})$, and by (6.25) we conclude $-\text{Curl } F = \Lambda_L^T$. The fact that F satisfies the boundary condition is straightforward, and thus $F \in \mathcal{F}_b^p$, achieving the proof. \square

REMARK 6.7. One cannot state that the above-solved variational problem is a complete solution to the dislocation problem, since what we have found in theorem 6.6 is merely a divergence-free deformation, which is presumably not the deformation associated with the dislocation problem in finite elasticity. Indeed, this result must be considered as a partial result towards this aim. One idea is to exploit the decomposition $F = Du + \text{Curl } V$ for the complete problem, and hence theorem 6.6 provides only the divergence-free part $\text{Curl } V$ of the true deformation tensor F . Solving the complete problem is an extremely difficult task, without additional assumptions on F or on the nature of the dislocations. To this respect some results are found in [37].

6.5.2. *A variational problem in finite gradient elasticity.* In this example, we treat a slightly different setting. We consider an energy involving F and the gradient of F , viz. $\mathcal{W} = \tilde{\mathcal{W}}(F, \nabla F)$ where there is an energetic term for the curl of F and another for its divergence. The term involving the divergence must be understood as a high-order perturbation, in the spirit of [18] (In their case, since no dislocations were considered, $\text{div } F = \Delta u$, with u the displacement was taken to the power 2 to account for the microstructure.), whereas the term related to the curl of F will

again be called the defect term, in view of the problem of the previous sections. Thus for some fixed $\epsilon > 0$, we consider

$$\tilde{\mathcal{W}}(F) := \mathcal{W}(F) + \epsilon \mathcal{W}_{\text{per}}(\text{div } F). \tag{6.30}$$

The same assumptions (6.20) and (6.21) hold for \mathcal{W} , and for the perturbation, one assumes that

$$\mathcal{W}_{\text{per}}(\text{div } F) \geq \gamma \|\text{div } F\|_{L^q}^q, \tag{6.31}$$

for $q > 3$, where 3 is precisely the space dimension. Note that such an assumption could not be made on the whole gradient of F , since we know that $\text{Curl } F$ is a measure concentrated in the dislocations. Indeed, notice that since $-\text{Curl } F = \Lambda_{\mathcal{L}}^T$, the divergence of F is not related to the dislocation density.

We assume that the map $\hat{F} \in \mathcal{A}^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with $-\text{Curl } \hat{F} = \Lambda_{\alpha}^T$ on $\hat{\Omega}$ satisfies the additional property that $\text{div } \hat{F} \in L^q(\hat{\Omega}, \mathbb{R}^3)$. Let $b \in 2\pi\mathbb{Z}^3$ a fixed Burgers vector, let $p, q > 1$, then we define the class of admissible functions as

$$\begin{aligned} \mathcal{F}_b^{p,q} := \{F \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}) : \text{div } F \in L^q(\hat{\Omega}, \mathbb{R}^3), -\text{Curl } F = b \otimes \mathcal{L} \\ \text{for some closed integral 1-current } \mathcal{L}, \text{ and } F = \hat{F} \text{ on } \hat{\Omega} \setminus \Omega\}. \end{aligned} \tag{6.32}$$

The existence of a minimizer of \mathcal{W} in $\mathcal{F}_b^{p,q}$ is provided by the following:

THEOREM 6.8. *Let $1 < p < 2$ and $q > 3$. Assume that \mathcal{W} satisfies the assumptions of theorem 6.6. If \mathcal{W}_{per} satisfies (6.31) and is lower semicontinuous with respect to the weak convergence in L^q , then there exists a minimizer $F \in \mathcal{F}_b^{p,q}$ of $\tilde{\mathcal{W}}$.*

Proof. We will apply the direct method. Let $\{F_k\}_{k>0}$ be a minimizing sequence. From the coerciveness (i) we see that there exist $F \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$, $A \in L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3})$, $D \in L^p(\hat{\Omega})$, and $R \in L^p(\hat{\Omega}, \mathbb{R}^3)$ such that (6.24a)–(6.24c) hold, with moreover,

$$\text{div } F_k \rightharpoonup R \quad \text{weakly in } L^q(\hat{\Omega}, \mathbb{R}^3). \tag{6.33a}$$

We find a measure $\Lambda \in \mathcal{M}_b(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ with $\Lambda_k \rightharpoonup \Lambda$ weakly* in $\mathcal{M}_b(\hat{\Omega}, \mathbb{R}^{3 \times 3})$, where we have set $\Lambda_k = \Lambda_{\mathcal{L}_k} = -(\text{Curl } F_k)^T$. Again, [37, lemma 7.5] implies that there exists a regular dislocation current \mathcal{L} such that $(\mathcal{L}_k)_i \rightharpoonup \mathcal{L}_i$ in $\mathcal{D}_1(\hat{\Omega})$ and $\Lambda = \Lambda_{\mathcal{L}} = \sum_{i=1}^3 \mathcal{L}_i \otimes e_i$. In order to prove the Theorem, we have to show that $\text{div } F = R$, $-\text{Curl } F = \Lambda_{\mathcal{L}}$, $A = \text{adj} F$, and $D = \det F$. The Helmholtz decomposition gives

$$F_k = Dw_k + \text{Curl } G_k, \tag{6.34}$$

with $w_k \in W^{1,p}(\hat{\Omega}, \mathbb{R}^3)$ satisfying $-\Delta w_k = -\text{div } F_k$ with $\partial_N w_k = F_k N = \hat{F} N$ on $\partial \hat{\Omega}$ (note that $F_k N$ and $\hat{F} N$ have a distributional meaning, the divergences of F_k and \hat{F} being in L^p , see [36]), and $G_k \in \tilde{\mathcal{V}}^p(\hat{\Omega})$. Since $\text{div } F_k \in L^q(\hat{\Omega}, \mathbb{R}^3)$, with $q > 3$, by the regularity theory of elliptic problems and the Sobolev embedding Theorem, we find that $w_k \in \mathcal{C}^1(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ and that the L^∞ norm of their gradients

are bounded by a constant,

$$\|Dw_k\|_{L^\infty} < C. \tag{6.35}$$

Moreover, we have, up to a subsequence that

$$w_k \rightharpoonup w \quad \text{weakly in } W^{1,q}(\hat{\Omega}, \mathbb{R}^3), \tag{6.36}$$

for some $w \in W^{1,q}(\hat{\Omega}, \mathbb{R}^3)$.

Let us set $V_k := \text{Curl } G_k$. Now $-\text{Curl } V_k = \Lambda_{\mathcal{L}_k}^T$, and theorem 6.4 provides functions $u_k \in \mathcal{A}_p(\hat{\Omega}, \mathbb{T}^3)$ and $v_k \in \mathcal{C}^1(\hat{\Omega}, \mathbb{T}^3)$ such that $\nabla u_k - \nabla v_k = V_k$ satisfying

$$\partial \mathcal{G}_{u_k}(\omega) = \mathcal{L}_k \wedge \vec{b}(\omega), \tag{6.37}$$

for all $\omega \in \mathcal{D}^3(\hat{\Omega} \times \mathbb{T}^3)$, and

$$\|Dv_k\|_{L^\infty} \leq C|b \otimes \mathcal{L}_k|(\hat{\Omega}). \tag{6.38}$$

Thanks to (6.24a), (6.36), and (6.38), we can assume that there exist $u \in W^{1,p}(\hat{\Omega}, \mathbb{T}^3)$ and $v \in W^{1,p}(\hat{\Omega}, \mathbb{R}^{3 \times 3})$ such that $u_k \rightarrow u$ and $v_k \rightarrow v$ strongly in $L^p(\hat{\Omega}, \mathbb{R}^3)$,

$$\nabla u_k \rightharpoonup \nabla u \quad \text{weakly in } L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}), \tag{6.39}$$

and

$$\nabla v_k \rightharpoonup \nabla v \quad \text{weakly in } L^p(\hat{\Omega}, \mathbb{R}^{3 \times 3}). \tag{6.40}$$

Thanks to estimates (6.35) and (6.38), theorem 4.6 applies providing

$$M(\partial \mathcal{G}_{w_k + u_k - v_k}) \leq C(1 + |\Lambda_{\mathcal{L}_k}|(\hat{\Omega}))|\Lambda_{\mathcal{L}_k}|(\hat{\Omega}) < C. \tag{6.41}$$

This allows us to apply theorem 2.3, obtaining (6.28) and (6.29). Now, arguing as in theorem 6.6 all other claims follow straightforwardly, achieving the proof. \square

REMARK 6.9. Once again, this is a partial solution to the complete dislocation problem. In particular, the growth assumption on the deformation divergence seems relatively strong, which we would like to weaken in future works more oriented towards specific dislocation modelling (see [40, 41]). Note that also the Γ -convergence result as $\epsilon \rightarrow 0$ is a difficult open problem.

6.6. Physical motivations of the Euclidean-valued case

We conclude with a quick remark on a possible reason in order to also consider vector-valued displacements in variational problems, without giving explicit examples.

Let us recall that the dislocated solid is separated into two elastic half-spaces joined by atomic-level forces across their common interface, known as the glide plane. In the applications as shown so far, the energy is made of two terms, the elastic part and the defect part which is assumed as concentrated in L . With a view to time evolution of dislocation, an important physical term should also be

added on the glide plane S , which is prescribed in single crystals at moderate temperature ranges, the so-called ‘misfit’ energy taking into account nonlinear atomic interactions, viz.,

$$\mathcal{W}_{\text{mis}} := \int_S W_{\text{mis}}(\delta(x))dS(x)$$

where $0 < \delta(x) < |b|$ and W_{mis} is the interplanar potential energy density. This problem was originally introduced and solved by Nabarro, and received considerable attention nowadays since it permits a reasonable alternative to atomic models.

Appendix A. Proof of Theorem 5.1

Proof. We will follow the lines of the proof of theorem 4.1. We consider a smooth tubular neighbourhood D_ϵ of the curve L , and a neighbourhood S_δ , of width 2δ , of the surface S . Then we consider the currents $\partial\mathcal{G}_{u^\epsilon, \delta}$ in $\Omega \times \mathbb{R}^3$, with $u^{\epsilon, \delta} := u|_{\Omega \setminus (D_\epsilon \cup S_\delta)}$ and let first $\delta \rightarrow 0$, and then $\epsilon \rightarrow 0$.

To simplify the notation, we denote by ∂D_ϵ only the part of the boundary of D_ϵ that does not belong to S_δ , and similarly ∂S_δ is the boundary of S_δ which does not belong to D_ϵ . Therefore, for all $\omega \in \mathcal{D}^2(\Omega \times \mathbb{R}^3)$ it holds

$$\begin{aligned} \partial\mathcal{G}_{u^\epsilon, \delta}(\omega) &= \int_{\partial D_\epsilon} \omega_{ij}(x, u(x)) \tilde{M}_i^j(D(\Phi_{\perp \partial D_\epsilon}(x))) d\mathcal{H}^2(x) \\ &\quad + \int_{\partial S_\delta} \omega_{ij}(x, u(x)) \tilde{M}_i^j(D(\Phi_{\perp \partial S_\delta}(x))) d\mathcal{H}^2, \end{aligned} \tag{A.1}$$

where $\Phi := Id \times u$. Let us study the second term of the last expression. Using the coordinates $\{\tau_1, \tau_2, N\}$ on S , with τ_1 and τ_2 a tangent basis to S and N its unit normal, this can be written as

$$\begin{aligned} &\int_{S \setminus D_\epsilon} \omega_{12}(x + \delta N, u(x + \delta N)) + \sum_{k=4}^6 \omega_{1k}(x + \delta N, u(x + \delta N)) \\ &\quad \times \frac{\partial u_{k-3}}{\partial \tau_2}(x + \delta N) d\mathcal{H}^2(x) \\ &\quad - \int_{S \setminus D_\epsilon} \sum_{k=4}^6 \omega_{2k}(x + \delta N, u(x + \delta N)) \frac{\partial u_{k-3}}{\partial \tau_1}(x + \delta N) d\mathcal{H}^2(x) \\ &\quad - \int_{S \setminus D_\epsilon} \omega_{12}(x - \delta N, u(x - \delta N)) - \sum_{k=4}^6 \omega_{1k}(x - \delta N, u(x - \delta N)) \\ &\quad \times \frac{\partial u_{k-3}}{\partial \tau_2}(x - \delta N) d\mathcal{H}^2(x) \\ &\quad + \int_{S \setminus D_\epsilon} \sum_{k=4}^6 \omega_{2k}(x - \delta N, u(x - \delta N)) \frac{\partial u_{k-3}}{\partial \tau_1}(x - \delta N) d\mathcal{H}^2(x). \end{aligned} \tag{A.2}$$

Using the fact that $d(\partial S_\delta, L) \geq \epsilon$, we see that such a current, thanks to (3.6), has uniformly bounded mass for all δ so that the currents $\mathcal{G}_{u^\epsilon, \delta}$ converge to $\mathcal{G}_{u^\epsilon, \delta=0}$ in

the sense of currents. Letting $\delta \rightarrow 0$ in (A.2), taking into account that $u_k = b_k \hat{u}$, we obtain exactly the second, third, and fourth terms in (5.3), with the only exception that we are integrating on $S \setminus D_\epsilon$, viz.,

$$\begin{aligned} \partial \mathcal{G}_{u^\epsilon, 0}(\omega) &= \int_{\partial D_\epsilon} \omega_{ij}(x, u(x)) \tilde{M}_i^j(D(\Phi_{\perp \partial D_\epsilon}(x))) d\mathcal{H}^2(x) \\ &+ \int_{S \setminus D_\epsilon} (\omega_{12}(x, u(x) + b) - \omega_{12}(x, u(x))) d\mathcal{H}^2(x) \\ &+ \sum_{k=4}^6 \int_{S \setminus D_\epsilon} b_{k-3} (\omega_{1k}(x, u(x) + b) - \omega_{1k}(x, u(x))) \frac{\partial \hat{u}}{\partial \tau_2} d\mathcal{H}^2(x) \\ &- \sum_{k=4}^6 \int_{S \setminus D_\epsilon} b_{k-3} (\omega_{2k}(x, u(x) + b) - \omega_{2k}(x, u(x))) \frac{\partial \hat{u}}{\partial \tau_1} d\mathcal{H}^2(x). \end{aligned} \tag{A.3}$$

Now, we can explicitly write the first integral as in (4.11), that is,

$$\begin{aligned} \int_{\partial D_\epsilon} \tilde{\omega}_{23}(x, u(x)) d\mathcal{H}^2 + \sum_{k=4}^6 \int_0^{2\pi} \int_0^l \epsilon \tilde{\omega}_{2k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) \frac{\partial u_{k-3}}{\partial \tau}(\epsilon, \theta, \tau) d\tau d\theta \\ - \sum_{k=4}^6 \int_0^l \int_0^{2\pi} \epsilon \tilde{\omega}_{3k}(\epsilon, \theta, \tau, u(\epsilon, \theta, \tau)) \frac{\partial u_{k-3}}{\partial \sigma}(\epsilon, \theta, \tau) d\theta d\tau, \end{aligned}$$

and following the same computations as in the proof of theorem 4.1, neglecting the vanishing terms, we obtain the following expression

$$\begin{aligned} - \sum_{k=4}^6 \int_0^l \omega_{3k}(\hat{\tau}, u^+(\tau)) u_{k-3}(\epsilon, \theta, \tau) \Big|_0^{\theta=2\pi} d\tau \\ + \sum_{k=4}^6 \int_0^l \int_0^{2\pi} \frac{d}{d\theta} \omega_{3k}(\hat{\tau}, u^+(\tau)) u_{k-3}(\epsilon, \theta, \tau) d\theta d\tau, \end{aligned} \tag{A.4}$$

with $\hat{\tau} = (0, 0, \tau)$. Letting ϵ go to 0 and integrating by parts in θ as in (4.14), we obtain

$$- \sum_{k=4}^6 \int_0^l \int_0^{2\pi} \omega_{3k}(\hat{\tau}, u^+(\tau) + \frac{b}{2\pi} \theta) \frac{b_{k-3}}{2\pi} d\theta d\tau, \tag{A.5}$$

where we have used that $\lim_{\epsilon \rightarrow 0} u(\epsilon, \theta, \tau) = u^+(\tau) = u^+(\tau) + ((b\theta)/(2\pi))$. This last expression is exactly the first term of (5.3). To see that the second, third, and fourth terms of (A.3) tend to the correspondent terms in (5.3), it suffices to observe that the total mass of the currents represented by them are bounded (uniformly with respect to ϵ) thanks to the estimates (ii) and (iii) of theorem 3.10 so that $\mathcal{G}_{u^\epsilon, 0}$ converges to \mathcal{G}_u in the sense of currents and we are done.

It remains to prove (5.4). We argue as in the proof of theorem 4.6. The previous computations applied to the function w gives rise to a formula like (A.3) for the part of the boundary on S plus the term containing the 2×2 determinants of Dw .

These terms converge to the first four lines of (5.4). As for the part of the boundary on L we have an approximation as in (4.18), where the first row can be treated as in (A.4). This gives rise to $\mathcal{L} \wedge \vec{b}_w(\omega)$. The other non-vanishing terms are given by the expression (4.19), with the same notation. In contrast, we have that ω is not periodic anymore. Then arguing as in (4.19) we find the final expression

$$\sum_{4 \leq i \neq j \leq 6} \int_0^l \int_0^{2\pi} \omega_{ij} \left(\hat{\tau}, \frac{b\theta}{2\pi} + w(\hat{\tau}) \right) \frac{\partial v_{j-3}}{\partial \tau}(\hat{\tau}) \frac{b_{i-3}}{2\pi} d\theta d\tau = \mathcal{C}_w(\omega),$$

with $\hat{\tau} = (0, 0, \tau)$ as expressed in the local basis. Now \mathcal{C}_w can be estimated as in theorem 4.6. The estimates (5.6) and (5.7) are straightforward consequences of (5.3) and (5.4). □

Acknowledgements

The financial support of the FP7-IDEAS-ERC-StG #256872 (EntroPhase) is gratefully acknowledged by the first author. The present paper also benefits from the support of the GNAMPA (Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni) of INdAM (Istituto Nazionale di Alta Matematica). Second author was supported by the FCT Starting Grant 'Mathematical theory of dislocations: geometry, analysis, and modelling' (IF/00734/2013).

References

- 1 A. Aftalion, E. Sandier and S. Serfaty. Pinning phenomena in the Ginzburg-Landau model of superconductivity. *J. Math. Pures Appl. (9)* **80** (2001), 339–372.
- 2 G. Alberti. On maps whose distributional Jacobian is a measure. *Real Anal. Exch.* **33** (2008), 153–162.
- 3 G. Alberti, S. Baldo and G. Orlandi. Functions with prescribed singularities. *J. Eur. Math. Soc. (JEMS)* **5** (2003), 275–311.
- 4 F. Almgren. Deformations and multiple-valued functions. Geometric measure theory and the calculus of variations. *Proc. Summer Inst., Arcata/Calif. 1984, Proc. Symp. Pure Math.* **44** (1986), 29–30.
- 5 F. J. Almgren and E. H. Lieb. Singularities of energy minimizing maps from the ball to the sphere: Examples, counterexamples, and bounds. *Ann. Math. (2)* **128** (1988), 483–530.
- 6 J. M. Ball. Convexity conditions and existence theorems in nonlinear elasticity. *Arch. Ration. Mech. Anal.* **63** (1977), 337–403.
- 7 F. Bethuel, G. Orlandi and D. Smets. Motion of concentration sets in Ginzburg-Landau equations. *Ann. Fac. Sci. Toulouse, Math. (6)* **13** (2004), 3–43.
- 8 Ph. Bouafia, T. D. Pauw and C. Wang. Multiple valued maps into a separable Hilbert space that almost minimize their p Dirichlet energy or are squeeze and squash stationary. *Calc. Var. Partial Differ. Equ.* **54** (2015), 2167–2196.
- 9 J. Bourgain, A. Brezis and C. Wang. *Comptes Rendus Mathématique. Calc. Var. Partial Differ. Equ.* **338** (2004), 539543.
- 10 Y. Brenier. Non relativistic strings may be approximated by relativistic strings. *Methods Appl. Anal.* **12** (2005), 153–167.
- 11 H. Brézis, J.-M. Coron and E. H. Lieb. Harmonic maps with defects. *Commun. Math. Phys.* **107** (1986), 649–705.
- 12 P. G. Ciarlet. Three-dimensional elasticity. *Mathematical elasticity: three-dimensional elasticity* (Amsterdam: North-Holland, 1994).
- 13 S. Conti, A. Garroni and S. Müller. Singular kernels, multiscale decomposition of microstructure, and dislocation models. *Arch. Ration. Mech. Anal.* **199** (2011), 779–819.

- 14 S. Conti, A. Garroni and A. Massaccesi. Modeling of dislocations and relaxation of functionals on 1-currents with discrete multiplicity. *Calc. Var. Partial Differ. Equ.* **54** (2015a), 1847–1874.
- 15 S. Conti, A. Garroni and M. Ortiz. The line-tension approximation as the dilute limit of linear-elastic dislocations. *Arch. Ration. Mech. Anal.* **218** (2015b), 699–755.
- 16 C. De Lellis, C.-R. Grisanti and P. Tilli. Regular selections for multiple-valued functions. *Ann. Mat. Pura Appl. (4)* **183** (2004), 79–95.
- 17 H. Federer. *Geometric measure theory* (Berlin, Heidelberg, New York: Springer-Verlag 1969).
- 18 G. A. Francfort and S. Müller. Combined effects of homogenization and singular perturbations in elasticity. *J. Reine Angew. Math.* **454** (1994), 1–35.
- 19 M. Giaquinta, G. Modica and J. Souček. *Cartesian currents in the calculus of variations I. Cartesian currents*, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. 37* (Berlin: Springer, 1998a).
- 20 M. Giaquinta, G. Modica and J. Souček. *Cartesian currents in the calculus of variations II. Variational integrals. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. 38* (Berlin: Springer, 1998b).
- 21 M. Giaquinta, P. M. Mariano, G. Modica and D. Mucci. Currents and curvature varifolds in continuum mechanics. In *Nonlinear partial differential equations and related topics. Dedicated to Nina U. Uraltseva on the occasion of her 75th birthday*, pp. 97–117 (Providence, RI: American Mathematical Society (AMS), 2010).
- 22 D. Gilbarg and N. S. Trudinger. Elliptic partial differential equations of second order. Reprint of the 1998 ed. *Classics in Mathematics* (Berlin: Springer, 2001).
- 23 D. Henao and S. Serfaty. Energy estimates and cavity interaction for a critical-exponent cavitation model. *Commun. Pure Appl. Math.* **66** (2013), 1028–1101.
- 24 T. Hochrainer. Moving dislocations in finite plasticity: a topological approach. *Z. Angew. Math. Mech.* **93** (2013), 252–268.
- 25 T. Hochrainer and M. Zaiser. Fundamentals of a continuum theory of dislocations. *Proceedings of the International conference on Statistical Mechanics of Plasticity and Related Instabilities* (2005).
- 26 R. L. Jerrard and H. M. Soner. Functions of bounded higher variation. *Indiana Univ. Math. J.* **51** (2002), 645677.
- 27 H. Kleinert. *Gauge fields in condensed matter*, vol. 1 (Singapore: World Scientific Publishing, 1989).
- 28 H. Kozono and T. Yanagisawa. L^r -variational inequality for vector fields and the Helmholtz-Weyl decomposition in bounded domains. *Indiana Univ. Math. J.* **58** (2009), 1853–1920.
- 29 S. G. Krantz and H. R. Parks. Geometric integration theory. *Cornerstones* (Boston: Birkhäuser, 2008).
- 30 D. Mucci. Fractures and vector valued maps. *Calc. Var. Partial Differ. Equ.* **22** (2005), 391–420.
- 31 D. Mucci. A structure property of ‘vertical’ integral currents, with an application to the distributional determinant. *Rev. Mat. Complut.* **28** (2015), 49–83.
- 32 S. Müller. Det=det. A remark on the distributional determinant. *C. R. Acad. Sci., Paris, Sér. I 311* **1** (1990), 13–17.
- 33 S. Müller and M. Palombaro. Existence of minimizers for a polyconvex energy in a crystal with dislocations. *Calc. Var.* **31** (2008), 473–482.
- 34 M. Ortiz. “Microstructure Development and Evolution in Plasticity” Vienna Summer School on Microstructures, Vienna, Austria, September 25–29, 2000 (<http://www.ortiz.caltech.edu/presentations/index.html>).
- 35 E. Sandier and S. Serfaty. A product-estimate for Ginzburg-Landau and corollaries. *J. Funct. Anal.* **211** (2004), 219–244.
- 36 R. Scala and N. Van Goethem. Constraint reaction and the Peach-Koehler force for dislocation networks. *Mathematics and Mechanics of Complex Systems* **4** (2016), 105–128.
- 37 R. Scala and N. Van Goethem. Currents and dislocations at the continuum scale. *Methods Appl. Anal.* **23** (2016), 1–34.

- 38 N. Van Goethem and F. Dupret. A distributional approach to the geometry of 2D dislocations at the continuum scale. *Ann. Univ. Ferrara* **58** (2012), 407–434.
- 39 N. Van Goethem. Incompatibility-governed singularities in linear elasticity with dislocations. *Math. Mech. Solids* **22** (2017), 1688–1695.
- 40 R. Scala and N. Van Goethem. Variational evolution of dislocations in single crystals. *J. Nonlinear Sci.* (2018). doi:10.1007/s00332-018-9488-4.
- 41 R. Scala and N. Van Goethem. A variational approach to single crystals with dislocations. *SIAM J. Math. Anal.* (2019). doi:10.1137/18M1166572.